

Trigonometry up to Transformations

MCQ Test, Set 01

Time: 30 minutes

Maximum marks: 25

Name: _____ Class: _____ Date: _____

- There are 25 questions. Each has exactly one correct answer: choose A, B, C or D.
- Each correct answer carries 1 mark. There is no negative marking.
- The key sheet and worked solutions are at the end of this test.

1. If $\sec \theta = \frac{13}{5}$ and θ is acute, then $\tan \theta =$

- A. $\frac{5}{13}$ B. $\frac{12}{5}$ C. $\frac{5}{12}$ D. $\frac{12}{13}$

2. The radian measure of 210° is

- A. $\frac{7\pi}{6}$ B. $\frac{5\pi}{6}$ C. $\frac{7\pi}{12}$ D. $\frac{4\pi}{3}$

3. If $\theta = 230^\circ$, which of the following is positive?

- A. $\cos \theta$ B. $\sin \theta$ C. $\sec \theta$ D. $\tan \theta$

4. $\sin 60^\circ \cos 30^\circ - \cos 60^\circ \sin 30^\circ =$

- A. 1 B. $\frac{\sqrt{3}}{2}$ C. $\frac{1}{2}$ D. 0

5. $\cot(180^\circ - \theta) =$

- A. $\tan \theta$ B. $\cot \theta$
C. $-\cot \theta$ D. $-\tan \theta$

6. $\cos 1290^\circ =$

- A. $\frac{1}{2}$ B. $-\frac{1}{2}$ C. $\frac{\sqrt{3}}{2}$ D. $-\frac{\sqrt{3}}{2}$

7. The range of $\sec x$ is

- A. $(-1, 1)$ B. $(-\infty, -1] \cup [1, \infty)$
C. R D. $[-1, 1]$

8. The correct order of $\sin 1$, $\sin 2$, $\sin 3$ is

- A. $\sin 2 > \sin 1 > \sin 3$ B. $\sin 3 > \sin 2 > \sin 1$
C. $\sin 2 > \sin 3 > \sin 1$ D. $\sin 1 > \sin 2 > \sin 3$

9. $\frac{\sin 37^\circ}{\cos 53^\circ} =$

- A. -1 B. 0
C. 1 D. $\tan 37^\circ$

10. $\sin^6 \theta + \cos^6 \theta + 3 \sin^2 \theta \cos^2 \theta =$

- A. 0 B. 2 C. 3 D. 1

11. The least value of $\cos^2 \theta + \sec^2 \theta$ is

A. 2

B. 1

C. 0

D. 4

12. $\sin 75^\circ =$

A. $\frac{\sqrt{3}-1}{2\sqrt{2}}$

B. $\frac{\sqrt{2}+1}{2}$

C. $\frac{\sqrt{3}+1}{2}$

D. $\frac{\sqrt{3}+1}{2\sqrt{2}}$

13. $\tan 75^\circ =$

A. $\sqrt{3} + 1$

B. $2 + \sqrt{3}$

C. $2 - \sqrt{3}$

D. $\frac{\sqrt{3}+1}{2}$

14. $\sin^2 \left(\frac{\pi}{8} + \frac{\theta}{2} \right) - \sin^2 \left(\frac{\pi}{8} - \frac{\theta}{2} \right) =$

A. $\frac{1}{\sqrt{2}} \cos \theta$

B. $\sin \theta$

C. $\frac{1}{\sqrt{2}} \sin \theta$

D. $\sqrt{2} \sin \theta$

15. In $\triangle ABC$, $\tan A + \tan B + \tan C =$

A. 0

B. 1

C. $\tan A \tan B \tan C$

D. $-\tan A \tan B \tan C$

16. $(1 + \tan 8^\circ)(1 + \tan 37^\circ) =$

A. 1

B. 0

C. 4

D. 2

17. $2 \sin 22\frac{1}{2}^\circ \cos 22\frac{1}{2}^\circ =$

A. $\frac{1}{\sqrt{2}}$

B. $\frac{1}{2}$

C. 1

D. $\frac{\sqrt{3}}{2}$

18. If $\sin A = \frac{1}{4}$, then $\sin 3A =$

A. $\frac{3}{4}$

B. $\frac{13}{16}$

C. $\frac{11}{16}$

D. $\frac{1}{2}$

19. $\sin 18^\circ =$

A. $\frac{\sqrt{10+2\sqrt{5}}}{4}$

B. $\frac{\sqrt{5}-1}{2}$

C. $\frac{\sqrt{5}+1}{4}$

D. $\frac{\sqrt{5}-1}{4}$

20. $\cot 15^\circ - \tan 15^\circ =$

A. $2\sqrt{3}$

B. 4

C. $\frac{2}{\sqrt{3}}$

D. $\sqrt{3}$

21. $\cos \frac{\pi}{7} + \cos \frac{3\pi}{7} + \cos \frac{5\pi}{7} =$

A. 1

B. $-\frac{1}{2}$

C. 0

D. $\frac{1}{2}$

22. $\sin 20^\circ \sin 40^\circ \sin 80^\circ =$

A. $\frac{1}{8}$

B. $\frac{\sqrt{3}}{8}$

C. $\frac{3}{8}$

D. $\frac{\sqrt{3}}{4}$

23. $\sqrt{1 + \sin 50^\circ} - \sqrt{1 - \sin 50^\circ} =$

A. $2 \sin 50^\circ$

B. $2 \sin 25^\circ$

C. $\sin 25^\circ$

D. $2 \cos 25^\circ$

24. $2 \sin 75^\circ \cos 15^\circ =$

A. $\frac{2+\sqrt{3}}{2}$

B. $\frac{\sqrt{3}}{2}$

C. $\frac{2-\sqrt{3}}{2}$

D. $\frac{1}{2}$

25. The period of $4 \cos(5x + 3) - 1$ is

A. $\frac{5\pi}{2}$

B. $\frac{2\pi}{5}$

C. 2π

D. $\frac{\pi}{5}$

Key sheet

1. B	2. A	3. D	4. C	5. C
6. D	7. B	8. A	9. C	10. D
11. A	12. D	13. B	14. C	15. C
16. D	17. A	18. C	19. D	20. A
21. D	22. B	23. B	24. A	25. B

Solutions

1. If $\sec \theta = \frac{13}{5}$ and θ is acute, then $\tan \theta =$
 $\sec \theta = \frac{\text{hyp}}{\text{adj}} = \frac{13}{5}$: take hyp = 13, adj = 5
 $\text{opp} = \sqrt{13^2 - 5^2} = \sqrt{144} = 12$
 $\tan \theta = \frac{\text{opp}}{\text{adj}} = \frac{12}{5}$
Ans: (B) $\frac{12}{5}$
-

2. The radian measure of 210° is
 $210^\circ = 210 \times \frac{\pi}{180} = \frac{7\pi}{6}$
Ans: (A) $\frac{7\pi}{6}$
-

3. If $\theta = 230^\circ$, which of the following is positive?
 $230^\circ \in \text{III}_Q$, where only tan, cot are +ve
Ans: (D) $\tan \theta$
-

4. $\sin 60^\circ \cos 30^\circ - \cos 60^\circ \sin 30^\circ =$
 $= \frac{\sqrt{3}}{2} \cdot \frac{\sqrt{3}}{2} - \frac{1}{2} \cdot \frac{1}{2} = \frac{3}{4} - \frac{1}{4}$
 $= \frac{1}{2}$
Ans: (C) $\frac{1}{2}$
-

5. $\cot(180^\circ - \theta) =$
 $180^\circ - \theta$ is of the form $180^\circ \pm \theta$ or $360^\circ \pm \theta$, so cot does not change
For acute θ , $180^\circ - \theta \in \text{II}_Q$, where cot is -ve
 $\therefore \cot(180^\circ - \theta) = -\cot \theta$
Ans: (C) $-\cot \theta$
-

6. $\cos 1290^\circ =$
 $\cos 1290^\circ = \cos(1080^\circ + 210^\circ) = \cos 210^\circ$
 $210^\circ \in \text{III}_Q$, where cos is -ve
 $\cos 210^\circ = \cos(180^\circ + 30^\circ) = -\cos 30^\circ = -\frac{\sqrt{3}}{2}$
Ans: (D) $-\frac{\sqrt{3}}{2}$
-

7. The range of $\sec x$ is
 $\sec x = \frac{1}{\cos x}$ and $-1 \leq \cos x \leq 1$, $\cos x \neq 0$
So $\sec x \leq -1$ or $\sec x \geq 1$
Ans: (B) $(-\infty, -1] \cup [1, \infty)$
-

8. The correct order of $\sin 1$, $\sin 2$, $\sin 3$ is
 $1 \approx 57^\circ$, $2 \approx 114^\circ$, $3 \approx 171^\circ$
 $\sin 2 = \sin 66^\circ$, $\sin 3 = \sin 9^\circ$
 $\sin x$ increases on $(0^\circ, 90^\circ)$: $\sin 66^\circ > \sin 57^\circ > \sin 9^\circ$
Ans: (A) $\sin 2 > \sin 1 > \sin 3$
-

9. $\frac{\sin 37^\circ}{\cos 53^\circ} =$

$$\cos 53^\circ = \cos(90^\circ - 37^\circ) = \sin 37^\circ$$

So the ratio is 1

Ans: (C) 1

10. $\sin^6 \theta + \cos^6 \theta + 3 \sin^2 \theta \cos^2 \theta =$

$$\sin^6 \theta + \cos^6 \theta = 1 - 3 \sin^2 \theta \cos^2 \theta$$

Adding $3 \sin^2 \theta \cos^2 \theta$ gives 1

Ans: (D) 1

11. The least value of $\cos^2 \theta + \sec^2 \theta$ is

Consider $\cos^2 \theta$ and $\sec^2 \theta$; both are +ve

$$A.M \geq G.M: \frac{\cos^2 \theta + \sec^2 \theta}{2} \geq \sqrt{\cos^2 \theta \cdot \sec^2 \theta} = \sqrt{1} = 1$$

$$\Rightarrow \cos^2 \theta + \sec^2 \theta \geq 2; \text{equality when } \cos^2 \theta = 1$$

Ans: (A) 2

12. $\sin 75^\circ =$

$$\sin(45^\circ + 30^\circ) = \sin 45^\circ \cos 30^\circ + \cos 45^\circ \sin 30^\circ$$

$$= \frac{1}{\sqrt{2}} \cdot \frac{\sqrt{3}}{2} + \frac{1}{\sqrt{2}} \cdot \frac{1}{2} = \frac{\sqrt{3}+1}{2\sqrt{2}}$$

Ans: (D) $\frac{\sqrt{3}+1}{2\sqrt{2}}$

13. $\tan 75^\circ =$

$$\tan(45^\circ + 30^\circ) = \frac{1 + \frac{1}{\sqrt{3}}}{1 - \frac{1}{\sqrt{3}}} = \frac{\sqrt{3}+1}{\sqrt{3}-1}$$

$$= \frac{(\sqrt{3}+1)^2}{2} = \frac{4+2\sqrt{3}}{2} = 2 + \sqrt{3}$$

Ans: (B) $2 + \sqrt{3}$

14. $\sin^2 \left(\frac{\pi}{8} + \frac{\theta}{2} \right) - \sin^2 \left(\frac{\pi}{8} - \frac{\theta}{2} \right) =$

$$= \sin \left(\frac{\pi}{4} \right) \sin \theta = \frac{1}{\sqrt{2}} \sin \theta$$

Ans: (C) $\frac{1}{\sqrt{2}} \sin \theta$

15. In $\triangle ABC$, $\tan A + \tan B + \tan C =$

$$A + B = 180^\circ - C \Rightarrow \frac{\tan A + \tan B}{1 - \tan A \tan B} = -\tan C$$

$$\Rightarrow \tan A + \tan B + \tan C = \tan A \tan B \tan C$$

Ans: (C) $\tan A \tan B \tan C$

16. $(1 + \tan 8^\circ)(1 + \tan 37^\circ) =$

$$8^\circ + 37^\circ = 45^\circ \Rightarrow (1 + \tan A)(1 + \tan B) = 2$$

Ans: (D) 2

17. $2 \sin 22\frac{1}{2}^\circ \cos 22\frac{1}{2}^\circ =$

$$2 \sin A \cos A = \sin 2A$$

$$= \sin 45^\circ = \frac{1}{\sqrt{2}}$$

Ans: (A) $\frac{1}{\sqrt{2}}$

18. If $\sin A = \frac{1}{4}$, then $\sin 3A =$

$$\sin 3A = 3 \sin A - 4 \sin^3 A = \frac{3}{4} - 4 \cdot \frac{1}{64} = \frac{3}{4} - \frac{1}{16} = \frac{11}{16}$$

Ans: (C) $\frac{11}{16}$

19. $\sin 18^\circ =$

Standard value: $\sin 18^\circ = \frac{\sqrt{5}-1}{4}$

Ans: (D) $\frac{\sqrt{5}-1}{4}$

20. $\cot 15^\circ - \tan 15^\circ =$

$$\cot A - \tan A = 2 \cot 2A$$

$$= 2 \cot 30^\circ = 2\sqrt{3}$$

Ans: (A) $2\sqrt{3}$

21. $\cos \frac{\pi}{7} + \cos \frac{3\pi}{7} + \cos \frac{5\pi}{7} =$

$$a_1 = \frac{\pi}{7}, d = \frac{2\pi}{7}, n = 3: \text{sum} = \frac{\cos \frac{3\pi}{7} \sin \frac{3\pi}{7}}{\sin \frac{\pi}{7}} = \frac{\sin \frac{6\pi}{7}}{2 \sin \frac{\pi}{7}} = \frac{1}{2}$$

Ans: (D) $\frac{1}{2}$

22. $\sin 20^\circ \sin 40^\circ \sin 80^\circ =$

$$\sin \theta \sin(60^\circ - \theta) \sin(60^\circ + \theta) = \frac{1}{4} \sin 3\theta \text{ with } \theta = 20^\circ$$

$$= \frac{1}{4} \sin 60^\circ = \frac{\sqrt{3}}{8}$$

Ans: (B) $\frac{\sqrt{3}}{8}$

23. $\sqrt{1 + \sin 50^\circ} - \sqrt{1 - \sin 50^\circ} =$

$$\sqrt{1 + \sin 50^\circ} = \cos 25^\circ + \sin 25^\circ, \sqrt{1 - \sin 50^\circ} = \cos 25^\circ - \sin 25^\circ \quad (25^\circ < 45^\circ)$$

$$\text{Difference} = 2 \sin 25^\circ$$

Ans: (B) $2 \sin 25^\circ$

24. $2 \sin 75^\circ \cos 15^\circ =$

$$2 \sin A \cos B = \sin(A + B) + \sin(A - B)$$

$$= \sin 90^\circ + \sin 60^\circ = 1 + \frac{\sqrt{3}}{2} = \frac{2+\sqrt{3}}{2}$$

Ans: (A) $\frac{2+\sqrt{3}}{2}$

25. The period of $4 \cos(5x + 3) - 1$ is

$$\text{Period of } \cos(ax + b) + k \text{ is } \frac{2\pi}{|a|} = \frac{2\pi}{5}$$

Ans: (B) $\frac{2\pi}{5}$
