

Trigonometry up to Transformations

MCQ Test, Set 08

Time: 30 minutes

Maximum marks: 25

Name: _____ Class: _____ Date: _____

- There are 25 questions. Each has exactly one correct answer: choose A, B, C or D.
- Each correct answer carries 1 mark. There is no negative marking.
- The key sheet and worked solutions are at the end of this test.

- $\frac{\tan \theta}{\sec \theta} =$
A. $\sin \theta$ **B.** $\cos \theta$ **C.** $\cot \theta$ **D.** $\csc \theta$
- $\frac{7\pi}{4}$ radians in degrees is
A. 225° **B.** 315° **C.** 135° **D.** 330°
- If $\tan \theta < 0$ and $\sec \theta > 0$, then θ lies in
A. I_Q **B.** II_Q **C.** III_Q **D.** IV_Q
- $4 \cos^2 45^\circ - \csc^2 45^\circ =$
A. 1 **B.** -2 **C.** 0 **D.** 2
- $\csc(270^\circ + \theta) =$
A. $-\csc \theta$ **B.** $\sec \theta$
C. $\csc \theta$ **D.** $-\sec \theta$
- $\sec \frac{23\pi}{3} =$
A. $\frac{2}{\sqrt{3}}$ **B.** $-\frac{2}{\sqrt{3}}$ **C.** 2 **D.** -2
- The function $\csc x$ is defined for all real x except
A. $x = 2n\pi, n \in \mathbb{Z}$ **B.** $x = n\pi, n \in \mathbb{Z}$
C. $x = (2n + 1)\frac{\pi}{2}, n \in \mathbb{Z}$ **D.** $x = \frac{n\pi}{4}, n \in \mathbb{Z}$
- If $x = \cos 3$ and $y = \cos 3^\circ$, then
A. $x = -y$ **B.** $x < y$ **C.** $x = y$ **D.** $x > y$
- $\tan 15^\circ \tan 35^\circ \tan 55^\circ \tan 75^\circ =$
A. 1 **B.** $\sqrt{3}$ **C.** 0 **D.** $\frac{1}{\sqrt{3}}$
- If $\sin \theta \cos \theta = \frac{3}{7}$, then $\sin^2 \theta + \cos^4 \theta =$
A. $\frac{31}{49}$ **B.** $\frac{9}{49}$ **C.** $\frac{22}{49}$ **D.** $\frac{40}{49}$

11. $\sqrt{1 - \sin^2 20^\circ} =$

A. $-\cos 20^\circ$

B. $\cos 20^\circ$

C. $-\sin 20^\circ$

D. $\sin 20^\circ$

12. $\cos 80^\circ \cos 20^\circ + \sin 80^\circ \sin 20^\circ =$

A. $\cos 100^\circ$

B. $-\frac{1}{2}$

C. $\frac{1}{2}$

D. $\frac{\sqrt{3}}{2}$

13. $\frac{1 - \tan 20^\circ}{1 + \tan 20^\circ} =$

A. $\tan 25^\circ$

B. $\tan 70^\circ$

C. $\tan 65^\circ$

D. $\tan 20^\circ$

14. $\cos^2(30^\circ + \theta) - \sin^2(30^\circ - \theta) =$

A. $\cos 2\theta$

B. $\frac{1}{2} \sin 2\theta$

C. $\frac{1}{2} \cos 2\theta$

D. $\frac{\sqrt{3}}{2} \cos 2\theta$

15. If $A + B + C = 90^\circ$, then $\cot A + \cot B + \cot C =$

A. $\cot A \cot B \cot C$

B. $-\cot A \cot B \cot C$

C. 0

D. 1

16. $(\cot 10^\circ - 1)(\cot 35^\circ - 1) =$

A. 1

B. 2

C. 0

D. -2

17. If $\sin A = \frac{2}{5}$, then $\cos 2A =$

A. $\frac{17}{25}$

B. $\frac{1}{5}$

C. $\frac{21}{25}$

D. $-\frac{17}{25}$

18. If $\cos A = \frac{1}{9}$ and A is acute, then $\sin \frac{A}{2} =$

A. $\frac{4}{9}$

B. $\frac{\sqrt{5}}{3}$

C. $\frac{2}{3}$

D. $\frac{1}{3}$

19. $\sin 18^\circ + \cos 36^\circ =$

A. 1

B. $\frac{1}{2}$

C. $\frac{\sqrt{5}}{2}$

D. $\frac{\sqrt{5}}{4}$

20. $\cos 10^\circ \cos 20^\circ \cos 40^\circ =$

A. $\frac{1}{8}$

B. $\frac{1}{8} \tan 10^\circ$

C. $\frac{1}{4} \cot 10^\circ$

D. $\frac{1}{8} \cot 10^\circ$

21. $1 + \cos \frac{2\pi}{7} + \cos \frac{4\pi}{7} + \cos \frac{6\pi}{7} =$

A. $\frac{1}{2}$

B. 1

C. $-\frac{1}{2}$

D. $\frac{3}{2}$

22. $\cot 80^\circ \cot 40^\circ + \cot 40^\circ \cot 20^\circ - \cot 20^\circ \cot 80^\circ =$

A. -3

B. 3

C. 1

D. $\sqrt{3}$

23. If $\frac{x}{y} = \frac{\sin 90^\circ}{\sin 30^\circ}$, then $\frac{x+y}{x-y} =$

A. $\frac{1}{3}$

B. $\sqrt{3}$

C. 2

D. 3

24. In $\triangle ABC$, $\sin 2A + \sin 2B + \sin 2C =$

A. $4 \sin A \sin B \cos C$

B. $1 + 4 \sin A \sin B \sin C$

C. $4 \cos A \cos B \cos C$

D. $4 \sin A \sin B \sin C$

25. The period of $\sin^5 x$ is

A. $\frac{2\pi}{5}$

B. π

C. 2π

D. 10π

Key sheet

1. A	2. B	3. D	4. C	5. D
6. C	7. B	8. B	9. A	10. D
11. B	12. C	13. A	14. C	15. A
16. B	17. A	18. C	19. C	20. D
21. A	22. B	23. D	24. D	25. C

Solutions

1. $\frac{\tan \theta}{\sec \theta} =$

$$\frac{\tan \theta}{\sec \theta} = \frac{\sin \theta}{\cos \theta} \times \cos \theta = \sin \theta$$

Ans: (A) $\sin \theta$

2. $\frac{7\pi}{4}$ radians in degrees is

$$\frac{7\pi}{4} = 7 \times 45^\circ = 315^\circ$$

Ans: (B) 315°

3. If $\tan \theta < 0$ and $\sec \theta > 0$, then θ lies in

$\tan$ is -ve and $\sec$ is +ve only in IV_Q

Ans: (D) IV_Q

4. $4 \cos^2 45^\circ - \csc^2 45^\circ =$

$$= 4 \left(\frac{1}{2}\right) - (\sqrt{2})^2 = 2 - 2$$

$$= 0$$

Ans: (C) 0

5. $\csc(270^\circ + \theta) =$

$270^\circ + \theta$ is of the form $90^\circ \pm \theta$ or $270^\circ \pm \theta$, so $\csc$ changes to $\sec$

For acute θ , $270^\circ + \theta \in IV_Q$, where $\csc$ is -ve

$$\therefore \csc(270^\circ + \theta) = -\sec \theta$$

Ans: (D) $-\sec \theta$

6. $\sec \frac{23\pi}{3} =$

$$\frac{23\pi}{3} = 23 \times 60^\circ = 1380^\circ$$

$$\sec 1380^\circ = \sec(1080^\circ + 300^\circ) = \sec 300^\circ$$

$300^\circ \in IV_Q$, where $\sec$ is +ve

$$\sec 300^\circ = \sec(360^\circ - 60^\circ) = +\sec 60^\circ = 2$$

Ans: (C) 2

7. The function $\csc x$ is defined for all real x except

$$\csc x = \frac{1}{\sin x} \text{ needs } \sin x \neq 0$$

$$\sin x = 0 \Rightarrow x = n\pi$$

Ans: (B) $x = n\pi, n \in \mathbb{Z}$

8. If $x = \cos 3$ and $y = \cos 3^\circ$, then

$$3 \approx 171^\circ \in II_Q, \text{ so } x = \cos 3 < 0$$

$$y = \cos 3^\circ > 0, \text{ so } x < y$$

Ans: (B) $x < y$

9. $\tan 15^\circ \tan 35^\circ \tan 55^\circ \tan 75^\circ =$

$\tan 15^\circ \tan 75^\circ = 1$ and $\tan 35^\circ \tan 55^\circ = 1$ (complementary pairs)

Ans: (A) 1

10. If $\sin \theta \cos \theta = \frac{3}{7}$, then $\sin^2 \theta + \cos^4 \theta =$

$\sin^2 \theta + \cos^4 \theta = \sin^2 \theta + \cos^2 \theta (1 - \sin^2 \theta) = 1 - \sin^2 \theta \cos^2 \theta$

$= 1 - \left(\frac{3}{7}\right)^2 = 1 - \frac{9}{49} = \frac{40}{49}$

Ans: (D) $\frac{40}{49}$

11. $\sqrt{1 - \sin^2 20^\circ} =$

$= \sqrt{\cos^2 20^\circ} = |\cos 20^\circ|$

$20^\circ \in I_Q \Rightarrow \cos 20^\circ > 0$, so $= \cos 20^\circ$

Ans: (B) $\cos 20^\circ$

12. $\cos 80^\circ \cos 20^\circ + \sin 80^\circ \sin 20^\circ =$

$= \cos(80^\circ - 20^\circ) = \cos 60^\circ = \frac{1}{2}$

Ans: (C) $\frac{1}{2}$

13. $\frac{1 - \tan 20^\circ}{1 + \tan 20^\circ} =$

$\frac{1 - \tan \theta}{1 + \tan \theta} = \tan(45^\circ - \theta)$

$= \tan 25^\circ$

Ans: (A) $\tan 25^\circ$

14. $\cos^2(30^\circ + \theta) - \sin^2(30^\circ - \theta) =$

$\cos^2 A - \sin^2 B = \cos(A + B) \cos(A - B)$

$= \cos 60^\circ \cos 2\theta = \frac{1}{2} \cos 2\theta$

Ans: (C) $\frac{1}{2} \cos 2\theta$

15. If $A + B + C = 90^\circ$, then $\cot A + \cot B + \cot C =$

$A + B = 90^\circ - C \Rightarrow \cot(A + B) = \tan C = \frac{1}{\cot C}$

$\frac{\cot A \cot B - 1}{\cot A + \cot B} = \frac{1}{\cot C} \Rightarrow \Sigma \cot A = \Pi \cot A$

Ans: (A) $\cot A \cot B \cot C$

16. $(\cot 10^\circ - 1)(\cot 35^\circ - 1) =$

$= (1 - \cot 10^\circ)(1 - \cot 35^\circ)$ and $10^\circ + 35^\circ = 45^\circ$

Value $= 2$

Ans: (B) 2

17. If $\sin A = \frac{2}{5}$, then $\cos 2A =$

$\cos 2A = 1 - 2 \sin^2 A = 1 - \frac{8}{25} = \frac{17}{25}$

Ans: (A) $\frac{17}{25}$

18. If $\cos A = \frac{1}{9}$ and A is acute, then $\sin \frac{A}{2} =$

$$\sin \frac{A}{2} = \sqrt{\frac{1 - \cos A}{2}} = \sqrt{\frac{1 - \frac{1}{9}}{2}} = \sqrt{\frac{8/9}{2}} = \sqrt{\frac{4}{9}} = \frac{2}{3}$$

Ans: (C) $\frac{2}{3}$

19. $\sin 18^\circ + \cos 36^\circ =$

$$= \frac{(\sqrt{5}-1)+(\sqrt{5}+1)}{4} = \frac{2\sqrt{5}}{4} = \frac{\sqrt{5}}{2}$$

Ans: (C) $\frac{\sqrt{5}}{2}$

20. $\cos 10^\circ \cos 20^\circ \cos 40^\circ =$

$$= \frac{\sin 80^\circ}{8 \sin 10^\circ} = \frac{\cos 10^\circ}{8 \sin 10^\circ} = \frac{1}{8} \cot 10^\circ$$

Ans: (D) $\frac{1}{8} \cot 10^\circ$

21. $1 + \cos \frac{2\pi}{7} + \cos \frac{4\pi}{7} + \cos \frac{6\pi}{7} =$

$$\cos \frac{2\pi}{7} + \cos \frac{4\pi}{7} + \cos \frac{6\pi}{7} = -\frac{1}{2}$$

$$\text{G.E} = 1 - \frac{1}{2} = \frac{1}{2}$$

Ans: (A) $\frac{1}{2}$

22. $\cot 80^\circ \cot 40^\circ + \cot 40^\circ \cot 20^\circ - \cot 20^\circ \cot 80^\circ =$

$$\cot(60^\circ + \theta) \cot(60^\circ - \theta) + \cot(60^\circ - \theta) \cot \theta - \cot \theta \cot(60^\circ + \theta) = 3 \text{ with } \theta = 20^\circ$$

Ans: (B) 3

23. If $\frac{x}{y} = \frac{\sin 90^\circ}{\sin 30^\circ}$, then $\frac{x+y}{x-y} =$

$$\frac{x+y}{x-y} = \frac{\tan 60^\circ}{\tan 30^\circ} = \frac{\sqrt{3}}{1/\sqrt{3}} = 3$$

Ans: (D) 3

24. In $\triangle ABC$, $\sin 2A + \sin 2B + \sin 2C =$

$$= 2 \sin(A+B) \cos(A-B) + 2 \sin C \cos C = 2 \sin C [\cos(A-B) - \cos(A+B)]$$

$$= 2 \sin C \cdot 2 \sin A \sin B = 4 \sin A \sin B \sin C$$

Ans: (D) $4 \sin A \sin B \sin C$

25. The period of $\sin^5 x$ is

$\sin^n x$ with n odd has period 2π

Ans: (C) 2π
