

Exponents (Level 2)**MCQ Test, Set 10**

Time: 30 minutes

Maximum marks: 25

Name: _____ Class: _____ Date: _____

- There are 25 questions. Each has exactly one correct answer: choose A, B, C or D.
- Each correct answer carries 1 mark. There is no negative marking.
- The key sheet and worked solutions are at the end of this test.

- $(\sqrt{3})^{-4} =$
A. $\frac{1}{9}$ **B.** 9 **C.** $\frac{1}{81}$ **D.** -6
- Which of these is undefined?
A. 5^0 **B.** 0^0 **C.** $(-5)^0$ **D.** 0^5
- $216^2 \times 36^{-2} =$
A. 6^{10} **B.** $\frac{1}{36}$ **C.** 36 **D.** 1
- $(36)^{\frac{3}{2}} \div (128)^{\frac{2}{7}} =$
A. $\frac{1}{54}$ **B.** 108 **C.** 3 **D.** 54
- $(-\frac{32}{243})^{-\frac{1}{5}} =$
A. $-\frac{2}{3}$ **B.** $-\frac{243}{32}$ **C.** $-\frac{3}{2}$ **D.** $\frac{3}{2}$
- $(\frac{1}{9})^{-\frac{1}{2}} - (\frac{1}{8})^{-\frac{1}{3}} =$
A. -1 **B.** $\frac{5}{2}$ **C.** $-\frac{1}{6}$ **D.** 1
- $\frac{4^{\frac{4}{5}} \times 6^{\frac{3}{5}}}{12^{\frac{3}{5}}} =$
A. $\frac{1}{2}$ **B.** 2 **C.** 4 **D.** 1
- $(x^{a-b})^{a+b} \div (x^{a-c})^{a+c} =$
A. $x^{b^2-c^2}$ **B.** $x^{2a^2-b^2-c^2}$ **C.** 1 **D.** $x^{c^2-b^2}$
- If $a = -2$, $b = -3$ and $c = 5$, then $a^3 + b^3 + c^3 =$
A. 90 **B.** -90 **C.** 0 **D.** 30
- If $\frac{7^{4x} \times 49^2 \times 343}{7^x} = 7^1$, then $x =$
A. $\frac{8}{3}$ **B.** -6 **C.** $-\frac{6}{5}$ **D.** -2
- If $49^x = (7\sqrt{7})^2$, then $x =$
A. $\frac{1}{2}$ **B.** 3 **C.** 1 **D.** $\frac{3}{2}$

12.	If $2^{x-3} \times 13^{x-4} = 52$, then $x =$			
	A. 5	B. -5	C. 6	D. 4
13.	If $7^{2y} = 25$, then $7^{-y} =$			
	A. $\frac{1}{5}$	B. -5	C. 5	D. $\frac{1}{25}$
14.	If $49^x = \frac{1}{343}$, then $x =$			
	A. $\frac{3}{2}$	B. $-\frac{3}{2}$	C. $-\frac{2}{3}$	D. -3
15.	If $4^x + 2^{2x+1} = 48$, then $x =$			
	A. 1	B. 2	C. 4	D. 3
16.	If $4^{x+1} = 64$, then $5^{x+1} =$			
	A. $\frac{1}{125}$	B. 625	C. 125	D. 25
17.	$\frac{1}{1+4^{\frac{1}{2}}} + \frac{1}{1+4^{-\frac{1}{2}}} =$			
	A. 2	B. 1	C. $\frac{2}{3}$	D. $\frac{1}{3}$
18.	$\left(\frac{x^{p^2+q^2}}{x^{pq}}\right)^{p+q} \cdot \left(\frac{x^{q^2+r^2}}{x^{qr}}\right)^{q+r} =$			
	A. $x^{p^3+r^3}$	B. $x^{p^3+2q^3+r^3}$	C. $x^{p^3+q^3+r^3}$	D. $x^{p^3-r^3}$
19.	If $2^p = 3$, $3^q = 5$, $5^r = 7$ and $7^t = \frac{1}{8}$, then $pqrt =$			
	A. 3	B. $\frac{1}{8}$	C. -3	D. 1
20.	If $7^x = 2^y = 56^z$, then $\frac{1}{z} =$			
	A. $\frac{1}{x} + \frac{3}{y}$	B. $\frac{3}{x} + \frac{1}{y}$	C. $\frac{1}{x} + \frac{1}{y}$	D. $\frac{3}{x+y}$
21.	$(0.04)^{\frac{3}{2}} =$			
	A. 0.008	B. 0.06	C. 0.08	D. 0.8
22.	$\left(\frac{\sqrt{10}}{3}\right)^4 \div \left(\frac{\sqrt{10}}{3}\right)^6 =$			
	A. $\frac{100}{9}$	B. $\frac{10}{9}$	C. $\frac{9}{10}$	D. $\frac{9}{100}$
23.	$(x^{\frac{1}{3}} + x^{-\frac{1}{3}})(x^{\frac{2}{3}} + x^{-\frac{2}{3}} - 1) =$			
	A. $x - \frac{1}{x}$	B. 1	C. $x + \frac{1}{x}$	D. $x^2 + \frac{1}{x^2}$
24.	$(9)^{0.75} \times (9)^{0.75} =$			
	A. 27	B. 81	C. 3	D. 9
25.	$\left[\left\{\left(\frac{1}{11}\right)^2\right\}^{-3}\right]^{\frac{1}{2}} =$			
	A. 121	B. 11	C. $\frac{1}{1331}$	D. 1331

Key sheet

1. A	2. B	3. C	4. D	5. C
6. D	7. B	8. D	9. A	10. D
11. D	12. A	13. A	14. B	15. B
16. C	17. B	18. B	19. C	20. A
21. A	22. C	23. C	24. A	25. D

Solutions

1. $(\sqrt{3})^{-4} =$

$$(\sqrt{3})^{-4} = \left(3^{\frac{1}{2}}\right)^{-4} = 3^{-2} = \frac{1}{9}$$

Ans: (A) $\frac{1}{9}$

2. Which of these is undefined?

$a^0 = 1$ only when $a \neq 0$; the notes say 0^0 is undefined

Ans: (B) 0^0

3. $216^2 \times 36^{-2} =$

$$= (6^3)^2 \times (6^2)^{-2}$$

$$= 6^6 \times 6^{-4}$$

$$= 6^{6-4} = 6^2$$

$$= 36$$

Ans: (C) 36

4. $(36)^{\frac{3}{2}} \div (128)^{\frac{2}{7}} =$

$$(6^2)^{\frac{3}{2}} \div (2^7)^{\frac{2}{7}}$$

$$= 6^3 \div 2^2$$

$$= 216 \div 4 = 54$$

Ans: (D) 54

5. $\left(-\frac{32}{243}\right)^{-\frac{1}{5}} =$

$$\left(-\frac{32}{243}\right)^{-\frac{1}{5}} = \left[\left(-\frac{2}{3}\right)^5\right]^{-\frac{1}{5}}$$

$$= \left(-\frac{2}{3}\right)^{-1} = \left(-\frac{3}{2}\right)^1$$

$$= -\frac{3}{2}$$

Ans: (C) $-\frac{3}{2}$

6. $\left(\frac{1}{9}\right)^{-\frac{1}{2}} - \left(\frac{1}{8}\right)^{-\frac{1}{3}} =$

$$\left(\frac{1}{9}\right)^{-\frac{1}{2}} = \left[\left(\frac{1}{3}\right)^2\right]^{-\frac{1}{2}} = 3$$

$$\left(\frac{1}{8}\right)^{-\frac{1}{3}} = \left[\left(\frac{1}{2}\right)^3\right]^{-\frac{1}{3}} = 2$$

$$3 - 2 = 1$$

Ans: (D) 1

$$\begin{aligned}
 7. \frac{4^{\frac{4}{5}} \times 6^{\frac{3}{5}}}{12^{\frac{3}{5}}} &= \\
 &= \frac{(2^2)^{\frac{4}{5}} \times (2 \times 3)^{\frac{3}{5}}}{(2^2 \times 3)^{\frac{3}{5}}} \\
 &= 2^{\frac{8}{5} + \frac{3}{5} - \frac{6}{5}} \times 3^{\frac{3}{5} - \frac{3}{5}} \\
 &= 2^1 \times 3^0 \\
 &= 2
 \end{aligned}$$

Ans: (B) 2

$$\begin{aligned}
 8. (x^{a-b})^{a+b} \div (x^{a-c})^{a+c} &= \\
 &= x^{a^2-b^2} \div x^{a^2-c^2} \\
 &= x^{a^2-b^2-a^2+c^2} = x^{c^2-b^2}
 \end{aligned}$$

Ans: (D) $x^{c^2-b^2}$

$$\begin{aligned}
 9. \text{ If } a = -2, b = -3 \text{ and } c = 5, \text{ then } a^3 + b^3 + c^3 &= \\
 a + b + c = -2 - 3 + 5 = 0 & \\
 \text{So } a^3 + b^3 + c^3 = 3abc = 3 \times (-2) \times (-3) \times 5 = 90 &
 \end{aligned}$$

Ans: (A) 90

$$\begin{aligned}
 10. \text{ If } \frac{7^{4x} \times 49^2 \times 343}{7^x} = 7^1, \text{ then } x &= \\
 \frac{7^{4x}}{7^x} \times (7^2)^2 \times 7^3 = 7^1 & \\
 \Rightarrow 7^{3x} \times 7^4 \times 7^3 = 7^1 & \\
 \Rightarrow 7^{3x+7} = 7^1 & \\
 \Rightarrow 3x + 7 = 1 \Rightarrow 3x = -6 \Rightarrow x = -2 &
 \end{aligned}$$

Ans: (D) -2

$$\begin{aligned}
 11. \text{ If } 49^x = (7\sqrt{7})^2, \text{ then } x &= \\
 7\sqrt{7} = 7^1 \times 7^{\frac{1}{2}} = 7^{\frac{3}{2}} & \\
 (7^2)^x = \left(7^{\frac{3}{2}}\right)^2 & \\
 \Rightarrow 7^{2x} = 7^3 & \\
 \Rightarrow 2x = 3 \Rightarrow x = \frac{3}{2} &
 \end{aligned}$$

Ans: (D) $\frac{3}{2}$

$$\begin{aligned}
 12. \text{ If } 2^{x-3} \times 13^{x-4} = 52, \text{ then } x &= \\
 52 = 2^2 \times 13^1 & \\
 \Rightarrow 2^{x-3} \times 13^{x-4} = 2^2 \times 13^1 & \\
 \Rightarrow x - 3 = 2 \text{ and } x - 4 = 1 & \\
 \Rightarrow x = 5 \text{ and } x = 5 & \\
 \therefore x = 5 &
 \end{aligned}$$

Ans: (A) 5

13. If $7^{2y} = 25$, then $7^{-y} =$

$$(7^y)^2 = 5^2$$

$$\Rightarrow 7^y = 5$$

$$7^{-y} = \frac{1}{7^y} = \frac{1}{5}$$

Ans: (A) $\frac{1}{5}$

14. If $49^x = \frac{1}{343}$, then $x =$

$$(7^2)^x = \frac{1}{7^3}$$

$$\Rightarrow 7^{2x} = 7^{-3}$$

$$\Rightarrow 2x = -3$$

$$\Rightarrow x = -\frac{3}{2}$$

Ans: (B) $-\frac{3}{2}$

15. If $4^x + 2^{2x+1} = 48$, then $x =$

$$(2^2)^x + 2^{2x+1} = 48$$

$$\Rightarrow 2^{2x} + 2^{2x+1} = 48$$

$$\Rightarrow 2^{2x} + 2 \times 2^{2x} = 48$$

$$\Rightarrow 2^{2x} (1 + 2) = 48$$

$$\Rightarrow 2^{2x} \times 3 = 48$$

$$\Rightarrow 2^{2x} = 48 \times \frac{1}{3} = 16 = 2^4$$

$$\Rightarrow 2x = 4 \Rightarrow x = 2$$

Ans: (B) 2

16. If $4^{x+1} = 64$, then $5^{x+1} =$

$$(2^2)^{x+1} = 2^6 \Rightarrow 2x + 2 = 6 \Rightarrow x = 2$$

$$5^{2+1} = 5^3 = 125$$

Ans: (C) 125

17. $\frac{1}{1+4^{\frac{1}{2}}} + \frac{1}{1+4^{-\frac{1}{2}}} =$

$$4^{\frac{1}{2}} = 2$$

$$\frac{1}{1+2} + \frac{1}{1+\frac{1}{2}} = \frac{1}{3} + \frac{2}{3} = 1$$

Ans: (B) 1

18. $\left(\frac{x^{p^2+q^2}}{x^{pq}}\right)^{p+q} \cdot \left(\frac{x^{q^2+r^2}}{x^{qr}}\right)^{q+r} =$
 $= x^{(p+q)(p^2-pq+q^2)} \cdot x^{(q+r)(q^2-qr+r^2)}$
 $= x^{p^3+q^3} \cdot x^{q^3+r^3} = x^{p^3+2q^3+r^3}$

Ans: (B) $x^{p^3+2q^3+r^3}$

19. If $2^p = 3$, $3^q = 5$, $5^r = 7$ and $7^t = \frac{1}{8}$, then $pqrt =$

$$5 = 3^q = (2^p)^q = 2^{pq}$$

$$7 = 5^r = (2^{pq})^r = 2^{pqr}$$

$$\frac{1}{8} = 7^t = (2^{pqr})^t = 2^{pqrt}$$

$$2^{pqrt} = \frac{1}{8} = 2^{-3}$$

$$\Rightarrow pqrt = -3$$

Ans: (C) -3

20. If $7^x = 2^y = 56^z$, then $\frac{1}{z} =$

$$\text{Let each} = k: 7 = k^{\frac{1}{x}}, 2 = k^{\frac{1}{y}}, 56 = k^{\frac{1}{z}}$$

$$56 = 7 \times 2^3, \text{ so } \frac{1}{z} = \frac{1}{x} + \frac{3}{y}$$

Ans: (A) $\frac{1}{x} + \frac{3}{y}$

21. $(0.04)^{\frac{3}{2}} =$

$$0.04 = \frac{4}{100} = \left(\frac{2}{10}\right)^2 = (0.2)^2$$

$$(0.04)^{\frac{3}{2}} = [(0.2)^2]^{\frac{3}{2}} = (0.2)^3 = 0.008$$

Ans: (A) 0.008

22. $\left(\frac{\sqrt{10}}{3}\right)^4 \div \left(\frac{\sqrt{10}}{3}\right)^6 =$

$$\frac{a^m}{a^n} = \frac{1}{a^{n-m}}$$

$$\left(\frac{\sqrt{10}}{3}\right)^4 \div \left(\frac{\sqrt{10}}{3}\right)^6 = \frac{1}{\left(\frac{\sqrt{10}}{3}\right)^2}$$

$$= \frac{9}{10}$$

Ans: (C) $\frac{9}{10}$

23. $(x^{\frac{1}{3}} + x^{-\frac{1}{3}})(x^{\frac{2}{3}} + x^{-\frac{2}{3}} - 1) =$

$$(a+b)(a^2+b^2-ab) = a^3+b^3 \text{ with } a = x^{\frac{1}{3}}, b = x^{-\frac{1}{3}}, ab = 1$$

$$= x + x^{-1} = x + \frac{1}{x}$$

Ans: (C) $x + \frac{1}{x}$

24. $(9)^{0.75} \times (9)^{0.75} =$

$$= (9)^{0.75+0.75} = (9)^{1.5}$$

$$1.5 = \frac{3}{2}$$

$$= (3^2)^{\frac{3}{2}} = 27$$

Ans: (A) 27

25. $\left[\left\{\left(\frac{1}{11}\right)^2\right\}^{-3}\right]^{\frac{1}{2}} =$

$$\frac{1}{11} = 11^{-1}$$

$$= 11^{(-1) \times 2 \times (-3) \times \frac{1}{2}}$$

$$= 11^3$$

$$= 1331$$

Ans: (D) 1331
