

Exponents (Level 2)

MCQ Test, Set 02

Time: 30 minutes

Maximum marks: 25

Name: _____ Class: _____ Date: _____

- There are 25 questions. Each has exactly one correct answer: choose A, B, C or D.
- Each correct answer carries 1 mark. There is no negative marking.
- The key sheet and worked solutions are at the end of this test.

1. $\sqrt[4]{5^3} =$

A. $5^{\frac{3}{4}}$

B. 5^{-1}

C. 5^{12}

D. $5^{\frac{4}{3}}$

2. If $a^5 = 3^{20}$ and $a > 0$, then $a =$

A. 81

B. 3^{100}

C. $3^{\frac{1}{4}}$

D. 3^{15}

3. $8^4 \times 2^{-11} =$

A. 8^{-7}

B. $\frac{1}{2}$

C. 2

D. 2^{23}

4. $(81)^{\frac{3}{4}} \div (32)^{\frac{3}{5}} =$

A. $\frac{8}{27}$

B. $\frac{27}{8}$

C. $\frac{27}{2}$

D. $\frac{3}{2}$

5. $(\sqrt[3]{27})^{-\frac{1}{2}} =$

A. $-\sqrt{3}$

B. $\sqrt{3}$

C. $\frac{1}{3}$

D. $\frac{1}{\sqrt{3}}$

6. $(\frac{16}{81})^{-\frac{3}{4}} =$

A. $-\frac{27}{8}$

B. $\frac{8}{27}$

C. $\frac{27}{8}$

D. $\frac{3}{2}$

7. $\frac{8^{\frac{1}{2}} \times 25^{-\frac{1}{4}}}{10^{-\frac{1}{2}}} =$

A. 4

B. 1

C. $\frac{1}{4}$

D. 8

8. $(\frac{5^4}{5^2})^3 \times (\frac{5^2}{5^3})^4 \times (\frac{5^3}{5^4})^2 =$

A. 1

B. 0

C. 5^{12}

D. 5

9. If $a + b + c = 0$ and $abc = 4$, then $a^3 + b^3 + c^3 =$

A. 0

B. 12

C. 4

D. 64

10. If $\frac{2^{4x} \times 16^2 \times 32}{2^x} = 2^7$, then $x =$

A. -6

B. $-\frac{6}{5}$

C. -2

D. $\frac{20}{3}$

11. If $25^x = (5\sqrt{5})^6$, then $x =$

A. 9

B. $\frac{9}{2}$

C. 3

D. $\frac{3}{2}$

12. If $3^{x-3} \times 2^{2x-4} = 48$, then $x =$
A. 4 **B.** 5 **C.** $\frac{5}{2}$ **D.** 7
-
13. If $3^x = 7$, then $9^x =$
A. 63 **B.** 14 **C.** $\frac{7}{9}$ **D.** 49
-
14. If $\left(\sqrt{\frac{3}{2}}\right)^{x-2} = \frac{4}{9}$, then $x =$
A. 0 **B.** 6 **C.** -2 **D.** -6
-
15. If $9^x + 3^{2x+1} = 324$, then $x =$
A. 2 **B.** 3 **C.** 4 **D.** 1
-
16. If $4^x = \frac{32}{2^x}$, then $x =$
A. $\frac{5}{2}$ **B.** $\frac{5}{3}$ **C.** $-\frac{5}{3}$ **D.** 5
-
17. $\frac{1}{1+x^{p-q}} + \frac{1}{1+x^{q-p}} =$
A. x^{p-q} **B.** 1 **C.** 0 **D.** 2
-
18. $\left(x^{a^2+ab+b^2}\right)^{a-b} =$
A. $x^{a^3+b^3}$ **B.** $x^{a^2-b^2}$ **C.** $x^{(a-b)^3}$ **D.** $x^{a^3-b^3}$
-
19. If $2^x = 7$, $7^y = 3$ and $3^z = 8$, then $xyz =$
A. -3 **B.** 3 **C.** 8 **D.** 1
-
20. If $2^x = 5^y = 10^{-z}$, then $\frac{1}{x} + \frac{1}{y} + \frac{1}{z} =$
A. -1 **B.** 2 **C.** 1 **D.** 0
-
21. $\sqrt[3]{125^{-2}} =$
A. -25 **B.** $\frac{1}{5}$ **C.** 25 **D.** $\frac{1}{25}$
-
22. $\left(\frac{2}{\sqrt{5}}\right)^7 \div \left(\frac{2}{\sqrt{5}}\right)^{10} =$
A. $\frac{125}{8}$ **B.** $\frac{8\sqrt{5}}{25}$ **C.** $\frac{5\sqrt{5}}{8}$ **D.** $\frac{8}{125}$
-
23. If $x = 2$, then $(x^{\frac{1}{3}} + x^{-\frac{1}{3}})(x^{\frac{2}{3}} + x^{-\frac{2}{3}} - 1) =$
A. 2 **B.** $\frac{17}{4}$ **C.** $\frac{5}{2}$ **D.** $\frac{3}{2}$
-
24. $(81)^{0.3} \times (81)^{0.45} =$
A. 3 **B.** $\frac{1}{27}$ **C.** 9 **D.** 27
-
25. $\left\{(2^3)^{-2}\right\}^{-\frac{1}{3}} =$
A. 64 **B.** 2 **C.** 4 **D.** $\frac{1}{4}$
-

Key sheet

1. A	2. A	3. C	4. B	5. D
6. C	7. A	8. A	9. B	10. C
11. B	12. A	13. D	14. C	15. A
16. B	17. B	18. D	19. B	20. D
21. D	22. C	23. C	24. D	25. C

Solutions

1. $\sqrt[4]{5^3} =$

$$\sqrt[n]{a^m} = a^{\frac{m}{n}}, \text{ so } \sqrt[4]{5^3} = 5^{\frac{3}{4}}$$

Ans: (A) $5^{\frac{3}{4}}$

2. If $a^5 = 3^{20}$ and $a > 0$, then $a =$

$$a^m = b^n \Rightarrow a = b^{\frac{n}{m}}$$

$$a = 3^{\frac{20}{5}} = 3^4 = 81$$

Ans: (A) 81

3. $8^4 \times 2^{-11} =$

$$= (2^3)^4 \times 2^{-11}$$

$$= 2^{12} \times 2^{-11}$$

$$= 2^{12-11} = 2^1$$

$$= 2$$

Ans: (C) 2

4. $(81)^{\frac{3}{4}} \div (32)^{\frac{3}{5}} =$

$$(3^4)^{\frac{3}{4}} \div (2^5)^{\frac{3}{5}}$$

$$= 3^3 \div 2^3$$

$$= 27 \div 8 = \frac{27}{8}$$

Ans: (B) $\frac{27}{8}$

5. $(\sqrt[3]{27})^{-\frac{1}{2}} =$

$$\sqrt[3]{27} = 3$$

$$(\sqrt[3]{27})^{-\frac{1}{2}} = 3^{-\frac{1}{2}}$$

$$= \frac{1}{3^{\frac{1}{2}}} = \frac{1}{\sqrt{3}}$$

Ans: (D) $\frac{1}{\sqrt{3}}$

6. $(\frac{16}{81})^{-\frac{3}{4}} =$

$$(\frac{16}{81})^{-\frac{3}{4}} = \left[\left(\frac{2}{3}\right)^4\right]^{-\frac{3}{4}}$$

$$= \left(\frac{2}{3}\right)^{-3} = \left(\frac{3}{2}\right)^3$$

$$= \frac{27}{8}$$

Ans: (C) $\frac{27}{8}$

$$\begin{aligned}
 7. \frac{8^{\frac{1}{2}} \times 25^{-\frac{1}{4}}}{10^{-\frac{1}{2}}} &= \\
 &= \frac{(2^3)^{\frac{1}{2}} \times (5^2)^{-\frac{1}{4}}}{(2 \times 5)^{-\frac{1}{2}}} \\
 &= 2^{\frac{3}{2} + \frac{1}{2}} \times 5^{-\frac{1}{2} + \frac{1}{2}} \\
 &= 2^2 \times 5^0 \\
 &= 4
 \end{aligned}$$

Ans: (A) 4

$$\begin{aligned}
 8. \left(\frac{5^4}{5^2}\right)^3 \times \left(\frac{5^2}{5^3}\right)^4 \times \left(\frac{5^3}{5^4}\right)^2 &= \\
 \text{Powers: } (4-2)(3) + (2-3)(4) + (3-4)(2) &= \\
 = 6 - 4 - 2 = 0 & \\
 \text{Value} = 5^0 = 1 &
 \end{aligned}$$

Ans: (A) 1

$$\begin{aligned}
 9. \text{ If } a + b + c = 0 \text{ and } abc = 4, \text{ then } a^3 + b^3 + c^3 &= \\
 a + b + c = 0 \Rightarrow a^3 + b^3 + c^3 = 3abc & \\
 = 3 \times 4 = 12 &
 \end{aligned}$$

Ans: (B) 12

$$\begin{aligned}
 10. \text{ If } \frac{2^{4x} \times 16^2 \times 32}{2^x} = 2^7, \text{ then } x &= \\
 \frac{2^{4x}}{2^x} \times (2^4)^2 \times 2^5 = 2^7 & \\
 \Rightarrow 2^{3x} \times 2^8 \times 2^5 = 2^7 & \\
 \Rightarrow 2^{3x+13} = 2^7 & \\
 \Rightarrow 3x + 13 = 7 \Rightarrow 3x = -6 \Rightarrow x = -2 &
 \end{aligned}$$

Ans: (C) -2

$$\begin{aligned}
 11. \text{ If } 25^x = (5\sqrt{5})^6, \text{ then } x &= \\
 5\sqrt{5} = 5^1 \times 5^{\frac{1}{2}} = 5^{\frac{3}{2}} & \\
 (5^2)^x = \left(5^{\frac{3}{2}}\right)^6 & \\
 \Rightarrow 5^{2x} = 5^9 & \\
 \Rightarrow 2x = 9 \Rightarrow x = \frac{9}{2} &
 \end{aligned}$$

Ans: (B) $\frac{9}{2}$

$$\begin{aligned}
 12. \text{ If } 3^{x-3} \times 2^{2x-4} = 48, \text{ then } x &= \\
 48 = 3^1 \times 2^4 & \\
 \Rightarrow 3^{x-3} \times 2^{2x-4} = 3^1 \times 2^4 & \\
 \Rightarrow x-3 = 1 \text{ and } 2x-4 = 4 & \\
 \Rightarrow x = 4 \text{ and } x = 4 & \\
 \therefore x = 4 &
 \end{aligned}$$

Ans: (A) 4

13. If $3^x = 7$, then $9^x =$

$$\begin{aligned} 9^x &= (3^2)^x = (3^x)^2 \\ &= 7^2 = 49 \end{aligned}$$

Ans: (D) 49

14. If $\left(\sqrt{\frac{3}{2}}\right)^{x-2} = \frac{4}{9}$, then $x =$

$$\begin{aligned} \left[\left(\frac{3}{2}\right)^{\frac{1}{2}}\right]^{x-2} &= \left(\frac{2}{3}\right)^2 \\ \Rightarrow \left(\frac{3}{2}\right)^{\frac{x-2}{2}} &= \left(\frac{3}{2}\right)^{-2} \\ \Rightarrow \frac{x-2}{2} &= -2 \\ \Rightarrow x-2 &= -4 \\ \Rightarrow x &= -2 \end{aligned}$$

Ans: (C) -2

15. If $9^x + 3^{2x+1} = 324$, then $x =$

$$\begin{aligned} (3^2)^x + 3^{2x+1} &= 324 \\ \Rightarrow 3^{2x} + 3^{2x+1} &= 324 \\ \Rightarrow 3^{2x} + 3 \times 3^{2x} &= 324 \\ \Rightarrow 3^{2x} (1 + 3) &= 324 \\ \Rightarrow 3^{2x} \times 4 &= 324 \\ \Rightarrow 3^{2x} &= 324 \times \frac{1}{4} = 81 = 3^4 \\ \Rightarrow 2x = 4 &\Rightarrow x = 2 \end{aligned}$$

Ans: (A) 2

16. If $4^x = \frac{32}{2^x}$, then $x =$

$$\begin{aligned} (2^2)^x \cdot 2^x &= 2^5 \Rightarrow 2^{3x} = 2^5 \\ \Rightarrow 3x = 5 &\Rightarrow x = \frac{5}{3} \end{aligned}$$

Ans: (B) $\frac{5}{3}$

17. $\frac{1}{1+x^p-q} + \frac{1}{1+x^q-p} =$

$$\begin{aligned} &= \frac{x^q}{x^q+x^p} + \frac{x^p}{x^p+x^q} \\ &= \frac{x^q+x^p}{x^p+x^q} = 1 \end{aligned}$$

Ans: (B) 1

18. $\left(x^{a^2+ab+b^2}\right)^{a-b} =$

$$\begin{aligned} &= x^{(a-b)(a^2+ab+b^2)} \\ &= x^{a^3+a^2b+ab^2-a^2b-ab^2-b^3} = x^{a^3-b^3} \end{aligned}$$

Ans: (D) $x^{a^3-b^3}$

19. If $2^x = 7$, $7^y = 3$ and $3^z = 8$, then $xyz =$

$$3 = 7^y = (2^x)^y = 2^{xy}$$

$$8 = 3^z = (2^{xy})^z = 2^{xyz}$$

$$2^{xyz} = 8 = 2^3$$

$$\Rightarrow xyz = 3$$

Ans: (B) 3

20. If $2^x = 5^y = 10^{-z}$, then $\frac{1}{x} + \frac{1}{y} + \frac{1}{z} =$

$$\text{Let each} = k: 2 = k^{\frac{1}{x}}, 5 = k^{\frac{1}{y}}, 10 = k^{-\frac{1}{z}}$$

$$10 = 2 \times 5, \text{ so } \frac{1}{x} + \frac{1}{y} = -\frac{1}{z}, \text{ so the sum is } 0$$

Ans: (D) 0

21. $\sqrt[3]{125^{-2}} =$

$$\sqrt[n]{a^m} = a^{\frac{m}{n}}$$

$$\sqrt[3]{125^{-2}} = 125^{-\frac{2}{3}} = (5^3)^{-\frac{2}{3}}$$

$$= 5^{-2} = \frac{1}{5^2} = \frac{1}{25}$$

Ans: (D) $\frac{1}{25}$

22. $\left(\frac{2}{\sqrt{5}}\right)^7 \div \left(\frac{2}{\sqrt{5}}\right)^{10} =$

$$\frac{a^m}{a^n} = \frac{1}{a^{n-m}}$$

$$\left(\frac{2}{\sqrt{5}}\right)^7 \div \left(\frac{2}{\sqrt{5}}\right)^{10} = \frac{1}{\left(\frac{2}{\sqrt{5}}\right)^3}$$

$$= \frac{5\sqrt{5}}{8}$$

Ans: (C) $\frac{5\sqrt{5}}{8}$

23. If $x = 2$, then $(x^{\frac{1}{3}} + x^{-\frac{1}{3}})(x^{\frac{2}{3}} + x^{-\frac{2}{3}} - 1) =$

$$(a + b)(a^2 + b^2 - ab) = a^3 + b^3 \text{ with } a = x^{\frac{1}{3}}, b = x^{-\frac{1}{3}}$$

$$ab = x^{\frac{1}{3}} \cdot x^{-\frac{1}{3}} = x^0 = 1$$

$$\text{So the value} = (x^{\frac{1}{3}})^3 + (x^{-\frac{1}{3}})^3 = x + \frac{1}{x}$$

$$= 2 + \frac{1}{2} = \frac{5}{2}$$

Ans: (C) $\frac{5}{2}$

24. $(81)^{0.3} \times (81)^{0.45} =$

$$= (81)^{0.3+0.45} = (81)^{0.75}$$

$$0.75 = \frac{3}{4}$$

$$= (3^4)^{\frac{3}{4}} = 27$$

Ans: (D) 27

25. $\left\{(2^3)^{-2}\right\}^{-\frac{1}{3}} =$

$$= 2^{3 \times (-2) \times (-\frac{1}{3})}$$

$$= 2^2$$

$$= 4$$

Ans: (C) 4
