

Exponents (Level 2)

MCQ Test, Set 09

Time: 30 minutes

Maximum marks: 25

Name: _____ Class: _____ Date: _____

- There are 25 questions. Each has exactly one correct answer: choose A, B, C or D.
- Each correct answer carries 1 mark. There is no negative marking.
- The key sheet and worked solutions are at the end of this test.

1. $\sqrt[6]{3^{12}} =$

A. 3^{72}

B. 729

C. 9

D. $\sqrt{3}$

2. If $a^4 = 9^6$ and $a > 0$, then $a =$

A. 3

B. $9^{\frac{2}{3}}$

C. 81

D. 27

3. $121^3 \times 11^{-4} =$

A. $\frac{1}{121}$

B. 11^{10}

C. 121

D. $\frac{1}{11}$

4. $(243)^{\frac{4}{5}} \times (4)^{\frac{3}{2}} =$

A. 6

B. 162

C. 648

D. $\frac{1}{648}$

5. $\left(\frac{1}{16}\right)^{-\frac{3}{4}} =$

A. 2

B. 8

C. -8

D. $\frac{1}{8}$

6. $\left(\frac{8}{343}\right)^{-\frac{2}{3}} =$

A. $-\frac{49}{4}$

B. $\frac{7}{2}$

C. $\frac{4}{49}$

D. $\frac{49}{4}$

7. $\frac{3^{\frac{4}{5}} \times 12^{\frac{1}{5}}}{8^{\frac{1}{5}}} =$

A. 6

B. 3

C. 2

D. 12

8. $\left(\frac{x^3}{x}\right)^4 \times \left(\frac{x}{x^2}\right)^3 \times \left(\frac{x^2}{x^3}\right)^5 =$

A. 0

B. x

C. x^{16}

D. 1

9. If $a = 1$, $b = 6$ and $c = -7$, then $a^3 + b^3 + c^3 =$

A. -42

B. -126

C. 0

D. 126

10. If $2^{4x} = 256$ and $10^y = 0.001$, then $x - y =$

A. -1

B. 5

C. 11

D. -5

11. If $2^{9x} \div 2^{4x} = \sqrt[4]{2^{30}}$, then $x =$

A. $\frac{15}{2}$

B. $\frac{15}{26}$

C. 6

D. $\frac{3}{2}$

12. If $11^{x-1} \times 2^x = 44$, then $x =$
A. 2 **B.** 1 **C.** 3 **D.** -2
-
13. If $5^x = 2$, then $25^x =$
A. 50 **B.** 27 **C.** $\frac{2}{25}$ **D.** 4
-
14. If $\left(\frac{5}{2}\right)^x \left(\frac{2}{5}\right)^{4x} = \frac{8}{125}$, then $x =$
A. 1 **B.** 3 **C.** $\frac{3}{5}$ **D.** -1
-
15. If $7^{x+1} - 7^x = 294$, then $x =$
A. -2 **B.** 2 **C.** 3 **D.** 1
-
16. If $16^x = \frac{8}{2^x}$, then $x =$
A. $\frac{3}{8}$ **B.** $\frac{3}{4}$ **C.** $\frac{3}{5}$ **D.** $-\frac{3}{5}$
-
17. $\frac{2}{1+x^{a-b}} + \frac{2}{1+x^{b-a}} =$
A. 0 **B.** 4 **C.** 2 **D.** 1
-
18. If $a = 2$ and $b = 1$, then $\left(\frac{2^{a^2+b^2}}{2^{ab}}\right)^{a+b} =$
A. 2^9 **B.** 2^7 **C.** 2^{27} **D.** 2^{15}
-
19. If $6^x = 5$, $5^y = 11$ and $11^z = 6$, then $xyz =$
A. -1 **B.** 6 **C.** 1 **D.** 0
-
20. If $a^x = b^y = c^z$ and $b = ac$, then $y =$
A. $\frac{xz}{x+z}$ **B.** $x + z$ **C.** $\frac{2xz}{x+z}$ **D.** $\frac{x+z}{xz}$
-
21. $(0.729)^{\frac{2}{3}} =$
A. 8.1 **B.** 0.81 **C.** 81 **D.** 0.486
-
22. $\left(\frac{4}{\sqrt{3}}\right)^3 \div \left(\frac{4}{\sqrt{3}}\right)^5 =$
A. $\frac{16}{3}$ **B.** $\frac{3}{16}$ **C.** $\frac{16}{9}$ **D.** $\frac{9}{16}$
-
23. $(x^{\frac{1}{3}} + y^{\frac{1}{3}})(x^{\frac{2}{3}} - x^{\frac{1}{3}}y^{\frac{1}{3}} + y^{\frac{2}{3}}) =$
A. $x + y$ **B.** $x^3 + y^3$ **C.** $x^2 + y^2$ **D.** $x - y$
-
24. If $\frac{x^{2.5}}{x^{0.5}} = 49$ and $x > 0$, then $x =$
A. $\sqrt{7}$ **B.** 49 **C.** 2401 **D.** 7
-
25. $\left\{(10^2)^{-3}\right\}^{\frac{1}{2}} =$
A. $\frac{1}{1000000}$ **B.** 1000 **C.** $\frac{1}{100}$ **D.** $\frac{1}{1000}$
-

Key sheet

1. C	2. D	3. C	4. C	5. B
6. D	7. A	8. D	9. B	10. B
11. D	12. A	13. D	14. A	15. B
16. C	17. C	18. A	19. C	20. A
21. B	22. B	23. A	24. D	25. D

Solutions

1. $\sqrt[6]{3^{12}} =$

$$\sqrt[6]{3^{12}} = 3^{\frac{12}{6}} = 3^2 = 9$$

Ans: (C) 9

2. If $a^4 = 9^6$ and $a > 0$, then $a =$

$$a = 9^{\frac{6}{4}} = 9^{\frac{3}{2}} = (3^2)^{\frac{3}{2}} = 3^3 = 27$$

Ans: (D) 27

3. $121^3 \times 11^{-4} =$

$$= (11^2)^3 \times 11^{-4}$$

$$= 11^6 \times 11^{-4}$$

$$= 11^{6-4} = 11^2$$

$$= 121$$

Ans: (C) 121

4. $(243)^{\frac{4}{5}} \times (4)^{\frac{3}{2}} =$

$$(3^5)^{\frac{4}{5}} \times (2^2)^{\frac{3}{2}}$$

$$= 3^4 \times 2^3$$

$$= 81 \times 8 = 648$$

Ans: (C) 648

5. $\left(\frac{1}{16}\right)^{-\frac{3}{4}} =$

$$\left(\frac{1}{16}\right)^{-\frac{3}{4}} = \left[\left(\frac{1}{2}\right)^4\right]^{-\frac{3}{4}}$$

$$= \left(\frac{1}{2}\right)^{-3} = (2)^3$$

$$= 8$$

Ans: (B) 8

6. $\left(\frac{8}{343}\right)^{-\frac{2}{3}} =$

$$\left(\frac{8}{343}\right)^{-\frac{2}{3}} = \left[\left(\frac{2}{7}\right)^3\right]^{-\frac{2}{3}}$$

$$= \left(\frac{2}{7}\right)^{-2} = \left(\frac{7}{2}\right)^2$$

$$= \frac{49}{4}$$

Ans: (D) $\frac{49}{4}$

7. $\frac{3^{\frac{4}{5}} \times 12^{\frac{1}{5}}}{8^{-\frac{1}{5}}} =$

$$= \frac{3^{\frac{4}{5}} \times (2^2 \times 3)^{\frac{1}{5}}}{(2^3)^{-\frac{1}{5}}}$$

$$= 2^{\frac{2}{5} + \frac{2}{5}} \times 3^{\frac{4}{5} + \frac{1}{5}}$$

$$= 2^1 \times 3^1$$

$$= 6$$

Ans: (A) 6

$$\begin{aligned}
 8. \left(\frac{x^3}{x}\right)^4 \times \left(\frac{x}{x^2}\right)^3 \times \left(\frac{x^2}{x^3}\right)^5 &= \\
 &= x^{2 \times 4} \times x^{-1 \times 3} \times x^{-1 \times 5} \\
 &= x^{8-3-5} = x^0 = 1
 \end{aligned}$$

Ans: (D) 1

$$\begin{aligned}
 9. \text{ If } a = 1, b = 6 \text{ and } c = -7, \text{ then } a^3 + b^3 + c^3 &= \\
 a + b + c = 1 + 6 - 7 = 0 & \\
 \text{So } a^3 + b^3 + c^3 = 3abc = 3 \times 1 \times 6 \times (-7) = -126 &
 \end{aligned}$$

Ans: (B) -126

$$\begin{aligned}
 10. \text{ If } 2^{4x} = 256 \text{ and } 10^y = 0.001, \text{ then } x - y &= \\
 2^{4x} = 256 = 2^8 \Rightarrow 4x = 8 \Rightarrow x = 2 & \\
 10^y = 0.001 = 10^{-3} \Rightarrow y = -3 & \\
 x - y = 2 - (-3) = 5 &
 \end{aligned}$$

Ans: (B) 5

$$\begin{aligned}
 11. \text{ If } 2^{9x} \div 2^{4x} = \sqrt[4]{2^{30}}, \text{ then } x &= \\
 \frac{2^{9x}}{2^{4x}} = 2^{\frac{30}{4}} & \\
 \Rightarrow 2^{9x-4x} = 2^{\frac{15}{2}} & \\
 \Rightarrow 5x = \frac{15}{2} \Rightarrow x = \frac{3}{2} &
 \end{aligned}$$

Ans: (D) $\frac{3}{2}$

$$\begin{aligned}
 12. \text{ If } 11^{x-1} \times 2^x = 44, \text{ then } x &= \\
 44 = 11^1 \times 2^2 & \\
 \Rightarrow 11^{x-1} \times 2^x = 11^1 \times 2^2 & \\
 \Rightarrow x - 1 = 1 \text{ and } x = 2 & \\
 \Rightarrow x = 2 \text{ and } x = 2 & \\
 \therefore x = 2 &
 \end{aligned}$$

Ans: (A) 2

$$\begin{aligned}
 13. \text{ If } 5^x = 2, \text{ then } 25^x &= \\
 25^x = (5^2)^x = (5^x)^2 & \\
 = 2^2 = 4 &
 \end{aligned}$$

Ans: (D) 4

$$\begin{aligned}
 14. \text{ If } \left(\frac{5}{2}\right)^x \left(\frac{2}{5}\right)^{4x} = \frac{8}{125}, \text{ then } x &= \\
 \left(\frac{5}{2}\right)^x = \left(\frac{2}{5}\right)^{-x} & \\
 \Rightarrow \left(\frac{2}{5}\right)^{-x} \cdot \left(\frac{2}{5}\right)^{4x} = \left(\frac{2}{5}\right)^3 & \\
 \Rightarrow \left(\frac{2}{5}\right)^{3x} = \left(\frac{2}{5}\right)^3 & \\
 \Rightarrow 3x = 3 \Rightarrow x = 1 &
 \end{aligned}$$

Ans: (A) 1

15. If $7^{x+1} - 7^x = 294$, then $x =$

$$7 \times 7^x - 7^x = 294$$

$$\Rightarrow 7^x (7 - 1) = 294$$

$$\Rightarrow 7^x \times 6 = 294$$

$$\Rightarrow 7^x = 294 \times \frac{1}{6} = 49 = 7^2$$

$$\Rightarrow x = 2$$

Ans: (B) 2

16. If $16^x = \frac{8}{2^x}$, then $x =$

$$(2^4)^x \cdot 2^x = 2^3 \Rightarrow 2^{5x} = 2^3$$

$$\Rightarrow 5x = 3 \Rightarrow x = \frac{3}{5}$$

Ans: (C) $\frac{3}{5}$

17. $\frac{2}{1+x^{a-b}} + \frac{2}{1+x^{b-a}} =$

$$= 2 \left(\frac{1}{1+x^{a-b}} + \frac{1}{1+x^{b-a}} \right) = 2 \times 1 = 2$$

Ans: (C) 2

18. If $a = 2$ and $b = 1$, then $\left(\frac{2^{a^2+b^2}}{2^{ab}} \right)^{a+b} =$

$$\left(\frac{2^{a^2+b^2}}{2^{ab}} \right)^{a+b} = 2^{(a+b)(a^2-ab+b^2)} = 2^{a^3+b^3}$$

$$= 2^{8+1} = 2^9$$

Ans: (A) 2^9

19. If $6^x = 5$, $5^y = 11$ and $11^z = 6$, then $xyz =$

$$11 = 5^y = (6^x)^y = 6^{xy}$$

$$6 = 11^z = (6^{xy})^z = 6^{xyz}$$

$$6^{xyz} = 6 = 6^1$$

$$\Rightarrow xyz = 1$$

Ans: (C) 1

20. If $a^x = b^y = c^z$ and $b = ac$, then $y =$

$$\text{Let each} = k: a = k^{\frac{1}{x}}, b = k^{\frac{1}{y}}, c = k^{\frac{1}{z}}$$

$$b = ac \Rightarrow k^{\frac{1}{y}} = k^{\frac{1}{x} + \frac{1}{z}} \Rightarrow \frac{1}{y} = \frac{x+z}{xz}$$

$$y = \frac{xz}{x+z}$$

Ans: (A) $\frac{xz}{x+z}$

21. $(0.729)^{\frac{2}{3}} =$

$$0.729 = \frac{729}{1000} = \left(\frac{9}{10} \right)^3 = (0.9)^3$$

$$(0.729)^{\frac{2}{3}} = [(0.9)^3]^{\frac{2}{3}} = (0.9)^2 = 0.81$$

Ans: (B) 0.81

22. $\left(\frac{4}{\sqrt{3}}\right)^3 \div \left(\frac{4}{\sqrt{3}}\right)^5 =$
 $\frac{a^m}{a^n} = \frac{1}{a^{n-m}}$
 $\left(\frac{4}{\sqrt{3}}\right)^3 \div \left(\frac{4}{\sqrt{3}}\right)^5 = \frac{1}{\left(\frac{4}{\sqrt{3}}\right)^2}$
 $= \frac{3}{16}$
Ans: (B) $\frac{3}{16}$

23. $(x^{\frac{1}{3}} + y^{\frac{1}{3}})(x^{\frac{2}{3}} - x^{\frac{1}{3}}y^{\frac{1}{3}} + y^{\frac{2}{3}}) =$
 $(a + b)(a^2 + b^2 - ab) = a^3 + b^3$ with $a = x^{\frac{1}{3}}, b = y^{\frac{1}{3}}$
 $= (x^{\frac{1}{3}})^3 + (y^{\frac{1}{3}})^3 = x + y$
Ans: (A) $x + y$

24. If $\frac{x^{2.5}}{x^{0.5}} = 49$ and $x > 0$, then $x =$
 $x^{2.5-0.5} = x^2 = 49$
 $x = 7$
Ans: (D) 7

25. $\left\{(10^2)^{-3}\right\}^{\frac{1}{2}} =$
 $= 10^{2 \times (-3) \times \frac{1}{2}}$
 $= 10^{-3}$
 $= \frac{1}{1000}$
Ans: (D) $\frac{1}{1000}$
