

Functions

MCQ Test, Set 04

Time: 30 minutes

Maximum marks: 25

Name: _____ Class: _____ Date: _____

1. There are 25 questions. Each has exactly one correct answer: choose A, B, C or D.
2. Each correct answer carries 1 mark. There is no negative marking.
3. The key sheet and worked solutions are at the end of this test.

1. If $f(x) = 5 - 3x$, then the image of 2 under f is
A. 1 **B.** -1 **C.** -11 **D.** 11
2. If $A = \{1, 2, 3\}$, $B = \{1, 2, 3, 4\}$ and $f : A \rightarrow B$ is defined by $f(x) = 5 - x$, then the range of f is
A. $\{1, 2, 3, 4\}$ **B.** $\{2, 3, 4\}$ **C.** $\{1, 2, 3\}$ **D.** $\{-4, -3, -2\}$
3. The degree of the polynomial function $f(x) = 8x^2 + 5x^4 - x + 6$ is
A. 4 **B.** 2 **C.** 8 **D.** 3
4. Which point lies on the graph of the identity function?
A. (0, 4) **B.** (4, 0) **C.** (-4, -4) **D.** (-4, 4)
5. The line $y = 7x - 14$ cuts the x-axis at
A. (0, -14) **B.** (-2, 0) **C.** (14, 0) **D.** (2, 0)
6. The discriminant of $2x^2 + 5x - 3 = 0$ is
A. 1 **B.** 29 **C.** 49 **D.** -1
7. If α, β are the roots of $2x^2 - 9x + 5 = 0$, then $\alpha + \beta =$
A. $\frac{9}{2}$ **B.** $-\frac{5}{2}$ **C.** $\frac{5}{2}$ **D.** $-\frac{9}{2}$
8. The range of $f(x) = x^2 - 4x + 1$, $x \in \mathbb{R}$, is
A. $[-3, \infty)$ **B.** $(-\infty, -3]$ **C.** $[2, \infty)$ **D.** $[1, \infty)$
9. The lowest point of the graph of $y = (x - 3)(x - 11)$ is
A. (-7, -16) **B.** (7, 33) **C.** (7, 16) **D.** (7, -16)
10. If $-2x^2 + 3x - 5 < 0$, then $x \in$
A. $\mathbb{R} - \{\frac{3}{4}\}$ **B.** ϕ
C. $\{\frac{3}{4}\}$ **D.** $\mathbb{R}$
11. If $25x^2 - 10x + 1 > 0$, then $x \in$
A. ϕ **B.** $\mathbb{R}$
C. $\mathbb{R} - \{\frac{1}{5}\}$ **D.** $\{\frac{1}{5}\}$

12. If $\log_5(x+2) = 2$, then $x =$
A. 10 **B.** 27 **C.** 23 **D.** 8
-
13. $\frac{1}{\log_4 2} =$
A. $\frac{1}{2}$ **B.** 2 **C.** 4 **D.** -2
-
14. Which of the following is negative?
A. $\log_5 1.2$ **B.** $\log_{1/4} 9$ **C.** $\log_{0.2} 0.5$ **D.** $\log_3 7$
-
15. $\log_7 2 \cdot \log_{13} 7 \cdot \log_8 13 =$
A. $\frac{2}{3}$ **B.** 3 **C.** $\frac{1}{3}$ **D.** $\frac{1}{2}$
-
16. If $|x| < x$, then $x \in$
A. ϕ **B.** $(-\infty, 0)$ **C.** $\mathbb{R}$ **D.** $(0, \infty)$
-
17. If $|x| \geq 6$, then $x \in$
A. $(-6, 6)$ **B.** $(-\infty, -6) \cup (6, \infty)$
C. $(-\infty, -6] \cup [6, \infty)$ **D.** $[-6, 6]$
-
18. If $|x-2| = |2x+1|$, then $x \in$
A. $\{-\frac{1}{3}, 3\}$ **B.** $\{-3, \frac{1}{3}\}$ **C.** $\{\frac{1}{3}\}$ **D.** $\{-3\}$
-
19. The solution set of $|2x-1| + x = 5$ is
A. $\{-4, 2\}$ **B.** $\{2\}$ **C.** $\{-2, 4\}$ **D.** $\{-4\}$
-
20. If $|x^2 + 2x| < 8$, then $x \in$
A. $(-2, 4)$ **B.** $[-4, 2]$ **C.** $\mathbb{R}$ **D.** $(-4, 2)$
-
21. The domain of $f(x) = \sqrt{x^2 - 1}$ is
A. $[-1, 1]$ **B.** $(-\infty, -1] \cup [1, \infty)$
C. $[1, \infty)$ **D.** $(-\infty, -1) \cup (1, \infty)$
-
22. The range of the step function $f(x) = [x]$ is
A. $\mathbb{N}$ **B.** $\mathbb{R}$ **C.** $[0, \infty)$ **D.** $\mathbb{Z}$
-
23. If $[x-1] < -3$, then $x \in$
A. $(-\infty, -1)$ **B.** $(-\infty, -2]$ **C.** $(-\infty, -3)$ **D.** $(-\infty, -2)$
-
24. $[-1.2] + \{-1.2\} =$
A. -0.8 **B.** 1.2 **C.** -1.2 **D.** -2.8
-
25. The domain of $f(x) = \log_2 x$ is
A. $(0, \infty)$ **B.** $(1, \infty)$ **C.** $\mathbb{R}$ **D.** $[0, \infty)$
-

Key sheet

1. B	2. B	3. A	4. C	5. D
6. C	7. A	8. A	9. D	10. D
11. C	12. C	13. B	14. B	15. C
16. A	17. C	18. B	19. A	20. D
21. B	22. D	23. D	24. C	25. A

Solutions

1. If $f(x) = 5 - 3x$, then the image of 2 under f is

$$f(2) = 5 - 3(2) = -1$$

Ans: (B) -1

2. If $A = \{1, 2, 3\}$, $B = \{1, 2, 3, 4\}$ and $f : A \rightarrow B$ is defined by $f(x) = 5 - x$, then the range of f is

$$f(1) = 5 - (1) = 4, f(2) = 5 - (2) = 3, f(3) = 5 - (3) = 2$$

$$\text{Range} = f(A) = \{2, 3, 4\} \text{ (not the codomain } B)$$

Ans: (B) $\{2, 3, 4\}$

3. The degree of the polynomial function $f(x) = 8x^2 + 5x^4 - x + 6$ is

$$\text{Degree} = \text{highest power of } x \text{ with non-zero coefficient} = 4$$

Ans: (A) 4

4. Which point lies on the graph of the identity function?

$$\text{On } y = x \text{ both coordinates are equal: } (-4, -4)$$

Ans: (C) $(-4, -4)$

5. The line $y = 7x - 14$ cuts the x-axis at

$$\text{put } y = 0: 7x - 14 = 0 \Rightarrow x = 2$$

Ans: (D) $(2, 0)$

6. The discriminant of $2x^2 + 5x - 3 = 0$ is

$$\Delta = b^2 - 4ac = 5^2 - 4(2)(-3) = 49$$

Ans: (C) 49

7. If α, β are the roots of $2x^2 - 9x + 5 = 0$, then $\alpha + \beta =$

$$a = 2, b = -9$$

$$\alpha + \beta = -\frac{b}{a} = \frac{9}{2}$$

Ans: (A) $\frac{9}{2}$

8. The range of $f(x) = x^2 - 4x + 1$, $x \in \mathbb{R}$, is

$$\text{Vertex} = \left(-\frac{b}{2a}, \frac{4ac-b^2}{4a}\right) = \left(2, \frac{4-16}{4}\right) = (2, -3)$$

$$\text{Range} = [-3, \infty)$$

Ans: (A) $[-3, \infty)$

9. The lowest point of the graph of $y = (x - 3)(x - 11)$ is

$$\text{It cuts the x-axis at } x = 3, 11; \text{ lowest point at } x = \frac{\alpha + \beta}{2} = 7$$

$$y = (7 - 3)(7 - 11) = -16$$

Ans: (D) $(7, -16)$

10. If $-2x^2 + 3x - 5 < 0$, then $x \in$
 $a = -2 < 0$ (downward parabola)
 $\Delta = b^2 - 4ac = 3^2 - 4(-2)(-5) = -31$
 $a < 0, \Delta < 0: y < 0$ for all $x \in \mathbb{R}$
 $\therefore x \in \mathbb{R}$
Ans: (D) $\mathbb{R}$

11. If $25x^2 - 10x + 1 > 0$, then $x \in$
 $a = 25 > 0$ (upward parabola)
 $\Delta = b^2 - 4ac = (-10)^2 - 4(25)(1) = 0$
 $\Delta = 0 \Rightarrow y = 25(x - \frac{1}{5})^2$
 $y > 0$ for all $x \neq \frac{1}{5}$, and $y = 0$ at $x = \frac{1}{5}$
 $\therefore x \in \mathbb{R} - \{\frac{1}{5}\}$
Ans: (C) $\mathbb{R} - \{\frac{1}{5}\}$

12. If $\log_5(x + 2) = 2$, then $x =$
 $x + 2 = 5^2 = 25 \Rightarrow x = 23$
Ans: (C) 23

13. $\frac{1}{\log_4 2} =$
 $\frac{1}{\log_4 2} = \log_2 4 = 2$
Ans: (B) 2

14. Which of the following is negative?
 $\log_a x < 0$ when $a > 1, 0 < x < 1$ or $0 < a < 1, x > 1$
 $\log_{1/4} 9$: base $< 1, 9 > 1$: negative
Ans: (B) $\log_{1/4} 9$

15. $\log_7 2 \cdot \log_{13} 7 \cdot \log_8 13 =$
 $= \log_8 2 = \log_{2^3} 2 = \frac{1}{3}$
Ans: (C) $\frac{1}{3}$

16. If $|x| < x$, then $x \in$
 $|x| \geq x$ for every real x
So $|x| < x$ has no solution: $x \in \phi$
Ans: (A) ϕ

17. If $|x| \geq 6$, then $x \in$
 $x \leq -6$ (or) $x \geq 6$
 $\Rightarrow x \in (-\infty, -6] \cup [6, \infty)$
Ans: (C) $(-\infty, -6] \cup [6, \infty)$

18. If $|x - 2| = |2x + 1|$, then $x \in$

$$x - 2 = \pm(2x + 1)$$

$$x - 2 = 2x + 1 \Rightarrow x + 3 = 0 \Rightarrow x = -3$$

$$x - 2 = -2x - 1 \Rightarrow 3x - 1 = 0 \Rightarrow x = \frac{1}{3}$$

Ans: (B) $\{-3, \frac{1}{3}\}$

19. The solution set of $|2x - 1| + x = 5$ is

Case (i) Suppose $x < \frac{1}{2}$ ($2x - 1 < 0$): $(1 - 2x) + x = 5 \Rightarrow 1 - x = 5 \Rightarrow x = -4$, which is in this case

Case (ii) Suppose $x \geq \frac{1}{2}$ ($2x - 1 \geq 0$): $(2x - 1) + x = 5 \Rightarrow 3x - 1 = 5 \Rightarrow x = 2$, which is in this case

So $x \in \{-4, 2\}$

Ans: (A) $\{-4, 2\}$

20. If $|x^2 + 2x| < 8$, then $x \in$

$$-8 < x^2 + 2x < 8$$

$$\Rightarrow x^2 + 2x + 8 > 0 \text{ and } x^2 + 2x - 8 < 0$$

$$x^2 + 2x + 8 > 0: a > 0, \Delta = -28 < 0 \Rightarrow x \in \mathbb{R}$$

$$x^2 + 2x - 8 < 0: \Delta = 36, x = \frac{-2 \pm 6}{2} = -4, 2 \Rightarrow x \in (-4, 2)$$

Common values: $x \in (-4, 2)$

Ans: (D) $(-4, 2)$

21. The domain of $f(x) = \sqrt{x^2 - 1}$ is

$\sqrt{x^2 - 1}$ is defined when $x^2 - 1 \geq 0$

$$\Rightarrow (x - 1)(x + 1) \geq 0$$

Domain = $(-\infty, -1] \cup [1, \infty)$

Ans: (B) $(-\infty, -1] \cup [1, \infty)$

22. The range of the step function $f(x) = [x]$ is

$[x]$ is always an integer and every integer n is $[n]$

Range = $\mathbb{Z}$

Ans: (D) $\mathbb{Z}$

23. If $[x - 1] < -3$, then $x \in$

$$[x - 1] = [x] - 1$$

$$[x] - 1 < -3 \Rightarrow [x] < -2 \Rightarrow [x] \leq -3$$

$$[x] = -3 \text{ for } -3 \leq x < -2: x \in (-\infty, -2)$$

Ans: (D) $(-\infty, -2)$

24. $[-1.2] + \{-1.2\} =$

$$[-1.2] = -2, \{-1.2\} = -1.2 - (-2) = 0.8$$

$$-2 + 0.8 = -1.2$$

(always $[x] + \{x\} = x$)

Ans: (C) -1.2

25. The domain of $f(x) = \log_2 x$ is

$\log_a N$ needs $N > 0$

Domain = $(0, \infty)$

Ans: (A) $(0, \infty)$
