

Functions

MCQ Test, Set 06

Time: 30 minutes

Maximum marks: 25

Name: _____ Class: _____ Date: _____

1. There are 25 questions. Each has exactly one correct answer: choose A, B, C or D.
2. Each correct answer carries 1 mark. There is no negative marking.
3. The key sheet and worked solutions are at the end of this test.

1. The pre-image of -8 under $f(x) = 3x + 4$ is
A. -4 **B.** -20 **C.** $-\frac{4}{3}$ **D.** 4
2. If $A = \{0, 1, 2\}$, $B = \{2, 5, 8, 11\}$ and $f : A \rightarrow B$ is defined by $f(x) = 3x + 2$, then the range of f is
A. $\{2, 5, 8\}$ **B.** $\{2, 5, 8, 11\}$ **C.** $\{0, 1, 2\}$ **D.** $\{-2, 1, 4\}$
3. The degree of the polynomial function $f(x) = x - 4x^3 + 9$ is
A. 1 **B.** 2 **C.** 3 **D.** 9
4. The identity function on $A = \{a, b, c\}$ is
A. $\{(a, a), (b, b), (c, c)\}$ **B.** $\{(a, b), (b, c), (c, a)\}$
C. $\{(a, c), (b, b), (c, a)\}$ **D.** $\{(a, a)\}$
5. The inclination of the line $y = 11x - 3$ with the x-axis is
A. greater than 90° **B.** less than 90°
C. equal to 90° **D.** equal to 0°
6. For $x^2 + 5x + 7 = 0$:
A. $\Delta = -23$, roots are not real (imaginary)
B. $\Delta = -3$, roots are real and distinct
C. $\Delta = -3$, roots are not real (imaginary)
D. $\Delta = 53$, roots are real and distinct
7. $x^2 - x - 12 =$
A. $(x - 3)(x - 4)$ **B.** $(x + 3)(x + 4)$ **C.** $(x - 3)(x + 4)$ **D.** $(x + 3)(x - 4)$
8. The maximum value of $-x^2 + 2x + 4$ is
A. 3 **B.** 4 **C.** 1 **D.** 5
9. The graph of $y = (x + 4)(x - 3)$ cuts the x-axis at
A. $(-4, 0)$ and $(-3, 0)$ **B.** $(-4, 0)$ and $(3, 0)$
C. $(4, 0)$ and $(-3, 0)$ **D.** $(4, 0)$ and $(3, 0)$

10. If $(5 - x)(x - 1) \leq 0$, then $x \in$
A. $(-\infty, 1] \cup [5, \infty)$ **B.** $(1, 5)$
C. $(-\infty, 1) \cup (5, \infty)$ **D.** $[1, 5]$
-
11. If $x^2 - 9 < 0$, then $x \in$
A. $(-\infty, -3] \cup [3, \infty)$ **B.** $[-3, 3]$
C. $(-3, 3)$ **D.** $(-\infty, -3) \cup (3, \infty)$
-
12. $2^7 = 128$ in logarithmic form is
A. $\log_7 128 = 2$ **B.** $\log_2 7 = 128$ **C.** $\log_2 128 = 7$ **D.** $\log_{128} 2 = 7$
-
13. $\log_3 54 - \log_3 2 =$
A. 3 **B.** 1 **C.** 27 **D.** $\log_3 52$
-
14. If $\log_3(x + 1) \geq 0$, then $x \in$
A. $[0, \infty)$ **B.** $(0, \infty)$ **C.** $(-1, 0]$ **D.** $[-1, \infty)$
-
15. If $\log_2(x^2 + 3x) - \log_2(x + 3) = 4$, then $x =$
A. 13 **B.** 4 **C.** 8 **D.** 16
-
16. If $x = 5$ and $y = -8$, then $|x + y| =$
A. 13 **B.** -3 **C.** 40 **D.** 3
-
17. If $|1 - x| > 4$, then $x \in$
A. $(-3, 5)$ **B.** $(-\infty, -3] \cup [5, \infty)$
C. $(-\infty, -3) \cup (5, \infty)$ **D.** $[-3, 5]$
-
18. If $|x^2 - 1| = 3$, then $x \in$
A. $\{-2, -\sqrt{2}, \sqrt{2}, 2\}$ **B.** $\{-2, 2\}$
C. $\{4\}$ **D.** $\{2\}$
-
19. The solution set of $|x + 1| + |x - 3| = 4$ is
A. ϕ **B.** $\{3\}$ **C.** $[-1, 3]$ **D.** $\mathbb{R}$
-
20. If $x^2 - 4|x| + 3 > 0$, then $x \in$
A. $(-1, 1)$ **B.** $(-\infty, -3) \cup (-1, 1) \cup (3, \infty)$
C. $(-\infty, 1) \cup (3, \infty)$ **D.** $(-\infty, -3) \cup (3, \infty)$
-
21. The domain of $f(x) = \frac{1}{\sqrt{x^2 - 5x + 4}}$ is
A. $(-\infty, 1] \cup [4, \infty)$ **B.** $(-\infty, 1) \cup (4, \infty)$
C. $(-\infty, -4) \cup (-1, \infty)$ **D.** $(1, 4)$
-
22. $[7.1] - [-7.1] =$
A. 0 **B.** -1 **C.** 14 **D.** 15
-

23. The domain of $f(x) = \sqrt{x - [x]}$ is

A. $[0, \infty)$

B. $\mathbb{R}$

C. $\mathbb{Z}$

D. ϕ

24. $\{3\} + \{-3\} =$

A. -6

B. 1

C. 0

D. 6

25. The range of $f(x) = x + \frac{1}{x}$ is

A. $\mathbb{R} - \{0\}$

B. $(-2, 2)$

C. $[2, \infty)$

D. $(-\infty, -2] \cup [2, \infty)$

Key sheet

1. A	2. A	3. C	4. A	5. B
6. C	7. D	8. D	9. B	10. A
11. C	12. C	13. A	14. A	15. D
16. D	17. C	18. B	19. C	20. B
21. B	22. D	23. B	24. C	25. D

Solutions

1. The pre-image of -8 under $f(x) = 3x + 4$ is

$$f(x) = -8 \Rightarrow 3x + 4 = -8 \Rightarrow x = -4$$

Ans: (A) -4

2. If $A = \{0, 1, 2\}$, $B = \{2, 5, 8, 11\}$ and $f : A \rightarrow B$ is defined by $f(x) = 3x + 2$, then the range of f is

$$f(0) = 3(0) + 2 = 2, f(1) = 3(1) + 2 = 5, f(2) = 3(2) + 2 = 8$$

$$\text{Range} = f(A) = \{2, 5, 8\} \text{ (not the codomain } B)$$

Ans: (A) $\{2, 5, 8\}$

3. The degree of the polynomial function $f(x) = x - 4x^3 + 9$ is

Degree = highest power of x with non-zero coefficient = 3

Ans: (C) 3

4. The identity function on $A = \{a, b, c\}$ is

$$I_A(x) = x \text{ for every } x \in A$$

$$\text{So } I_A = \{(a, a), (b, b), (c, c)\}$$

Ans: (A) $\{(a, a), (b, b), (c, c)\}$

5. The inclination of the line $y = 11x - 3$ with the x-axis is

$$\text{Here } a = 11 > 0$$

So the inclination is less than 90°

Ans: (B) less than 90°

6. For $x^2 + 5x + 7 = 0$:

$$\Delta = b^2 - 4ac = 5^2 - 4(1)(7) = -3$$

So the roots are not real (imaginary)

Ans: (C) $\Delta = -3$, roots are not real (imaginary)

7. $x^2 - x - 12 =$

$$x = \frac{1 \pm 7}{2} = -3, 4$$

$$x^2 - x - 12 = (x - \alpha)(x - \beta) = (x + 3)(x - 4)$$

Ans: (D) $(x + 3)(x - 4)$

8. The maximum value of $-x^2 + 2x + 4$ is

$$\text{Vertex} = \left(-\frac{b}{2a}, \frac{4ac - b^2}{4a}\right) = \left(1, \frac{-16 - 4}{-4}\right) = (1, 5)$$

$a < 0$: maximum value = 5

Ans: (D) 5

9. The graph of $y = (x + 4)(x - 3)$ cuts the x-axis at

$$\text{put } y = 0: (x + 4)(x - 3) = 0 \Rightarrow x = -4, 3$$

Ans: (B) $(-4, 0)$ and $(3, 0)$

10. If $(5 - x)(x - 1) \leq 0$, then $x \in$
 $y = (5 - x)(x - 1) = -x^2 + 6x - 5$
 $a = -1 < 0$ (downward parabola)
 $\Delta = b^2 - 4ac = 6^2 - 4(-1)(-5) = 16$
 $y = 0 \Rightarrow x = \frac{-6 \pm 4}{-2} = 1, 5$
Sign: $-$ outside 1 and 5, $+$ between them
 $\therefore x \in (-\infty, 1] \cup [5, \infty)$
Ans: (A) $(-\infty, 1] \cup [5, \infty)$

11. If $x^2 - 9 < 0$, then $x \in$
 $a = 1 > 0$ (upward parabola)
 $\Delta = b^2 - 4ac = 0^2 - 4(1)(-9) = 36$
 $y = 0 \Rightarrow x = \pm\sqrt{9} = -3, 3$
Sign: $+$ outside -3 and 3 , $-$ between them
 $\therefore x \in (-3, 3)$
Ans: (C) $(-3, 3)$

12. $2^7 = 128$ in logarithmic form is
 $a^x = N \Leftrightarrow x = \log_a N$
Base 2, power 7: $\log_2 128 = 7$
Ans: (C) $\log_2 128 = 7$

13. $\log_3 54 - \log_3 2 =$
 $\log_3 \frac{54}{2} = \log_3 27 = 3$
Ans: (A) 3

14. If $\log_3(x + 1) \geq 0$, then $x \in$
Base $3 > 1$: $\log_a T \geq 0 \Leftrightarrow T \geq 1$
 $x + 1 \geq 1 \Rightarrow x \geq 0$
Ans: (A) $[0, \infty)$

15. If $\log_2(x^2 + 3x) - \log_2(x + 3) = 4$, then $x =$
 $\log_2 \frac{x(x+3)}{x+3} = 4 \Rightarrow \log_2 x = 4 \Rightarrow x = 2^4 = 16$
Ans: (D) 16

16. If $x = 5$ and $y = -8$, then $|x + y| =$
 $|5 + (-8)| = |-3| = 3$
Ans: (D) 3

17. If $|1 - x| > 4$, then $x \in$

$$1 - x < -4 \text{ (or) } 1 - x > 4$$

$$\Rightarrow x < -3 \text{ (or) } x > 5 \text{ (dividing by a negative number turns the signs)}$$

$$\Rightarrow x \in (-\infty, -3) \cup (5, \infty)$$

$$\text{Ans: (C) } (-\infty, -3) \cup (5, \infty)$$

18. If $|x^2 - 1| = 3$, then $x \in$

$$x^2 - 1 = \pm 3$$

$$x^2 - 1 = 3 \Rightarrow x^2 - 4 = 0 \Rightarrow x = \pm\sqrt{4} = -2, 2$$

$$x^2 - 1 = -3 \Rightarrow x^2 + 2 = 0, \Delta = -8 < 0: \text{ no real } x$$

$$\text{Ans: (B) } \{-2, 2\}$$

19. The solution set of $|x + 1| + |x - 3| = 4$ is

Case (i) Suppose $x < -1$ ($x + 1 < 0, x - 3 < 0$): $(-x - 1) + (3 - x) = 4 \Rightarrow 2 - 2x = 4 \Rightarrow x = -1$, not in this case: no solution

Case (ii) Suppose $-1 \leq x \leq 3$ ($x + 1 \geq 0, x - 3 \leq 0$): $(x + 1) + (3 - x) = 4 \Rightarrow 4 = 4$: true for every x in this case

Case (iii) Suppose $x > 3$ ($x + 1 > 0, x - 3 > 0$): $(x + 1) + (x - 3) = 4 \Rightarrow 2x - 2 = 4 \Rightarrow x = 3$, not in this case: no solution

$$\text{So } x \in [-1, 3]$$

$$\text{Ans: (C) } [-1, 3]$$

20. If $x^2 - 4|x| + 3 > 0$, then $x \in$

$x = 0$ is the critical point

Case (i) Suppose $x < 0$: $|x| = -x, x^2 + 4x + 3 > 0 \Rightarrow (x + 1)(x + 3) > 0 \Rightarrow x \in (-\infty, -3) \cup (-1, \infty)$; common values $(-\infty, -3) \cup (-1, 0)$

Case (ii) Suppose $x \geq 0$: $|x| = x, x^2 - 4x + 3 > 0 \Rightarrow (x - 3)(x - 1) > 0 \Rightarrow x \in (-\infty, 1) \cup (3, \infty)$; common values $[0, 1) \cup (3, \infty)$

$$\text{From (i), (ii): } x \in (-\infty, -3) \cup (-1, 1) \cup (3, \infty)$$

$$\text{Ans: (B) } (-\infty, -3) \cup (-1, 1) \cup (3, \infty)$$

21. The domain of $f(x) = \frac{1}{\sqrt{x^2 - 5x + 4}}$ is

$$\frac{1}{\sqrt{x^2 - 5x + 4}} \text{ is defined when } x^2 - 5x + 4 > 0$$

$$\Rightarrow (x - 4)(x - 1) > 0$$

$$\text{Domain} = (-\infty, 1) \cup (4, \infty)$$

$$\text{Ans: (B) } (-\infty, 1) \cup (4, \infty)$$

22. $[7.1] - [-7.1] =$

$$[7.1] = 7, [-7.1] = -8$$

$$7 - (-8) = 15$$

$$\text{Ans: (D) } 15$$

23. The domain of $f(x) = \sqrt{x - [x]}$ is

Need $x - [x] \geq 0$, i.e. $[x] \leq x$

This is true for every real x : domain = $\mathbb{R}$

Ans: (B) $\mathbb{R}$

24. $\{3\} + \{-3\} =$

$\{3\} = 3 - 3 = 0$, $\{-3\} = -3 - (-3) = 0$

Ans: (C) 0

25. The range of $f(x) = x + \frac{1}{x}$ is

Minimum 2 at $x = 1$ (for $x > 0$), maximum -2 at $x = -1$ (for $x < 0$)

Range = $(-\infty, -2] \cup [2, \infty)$

Ans: (D) $(-\infty, -2] \cup [2, \infty)$
