EXPRESSWALA ACADEMY

Maths 1A: Algebra of Matrices

Question Paper, Set 08

Time: 1 hour 30 minutes | Maximum marks: 50

Name: _____ Class: _____ Date: _____

1. Answer all 20 questions. The marks for each question are shown in brackets.
2. Show your working for the 2, 3 and 5 mark questions.
3. The key sheet and worked solutions are at the end of this paper.

1. For $A = \begin{bmatrix} 0 & 1 \\ -8 & 0 \\ 6 & -5 \\ -9 & 8 \end{bmatrix}$, find the number of elements of A and $a_{41} + a_{22}$. [1 mark]

2. Write the matrix $A = [a_{ij}]_{2\times 3}$ whose elements are $a_{ij} = |i - 4j|$. [2 marks]

3. Name the type of the square matrix $\begin{bmatrix} -5 & 0 & 0 \\ 0 & -5 & 0 \\ 0 & 0 & -5 \end{bmatrix}$ (the most special type that fits). [1 mark]

4. If $\operatorname{tr}(A) = 1$ and $\operatorname{tr}(B) = 1$, find $\operatorname{tr}(4A + 2B)$. [1 mark]

5. If $\begin{bmatrix} 3x - y & 6 \\ 9 & x - y \end{bmatrix} = \begin{bmatrix} 17 & 6 \\ 9 & 3 \end{bmatrix}$, find the values of x and y. [2 marks]

6. Find the matrix X if $2X + \begin{bmatrix} 3 & -4 & -6 \\ 5 & 7 & 3 \end{bmatrix} = \begin{bmatrix} 3 & 2 & -4 \\ 3 & 3 & -5 \end{bmatrix}$. [2 marks]

7. Find X and Y if $2X + Y = \begin{bmatrix} -1 & 11 \\ 0 & 3 \end{bmatrix}$ and $X + 2Y = \begin{bmatrix} -2 & 7 \\ 0 & -6 \end{bmatrix}$. [3 marks]

8. If $A = \begin{bmatrix} 2 & 4 & 4 \end{bmatrix}$ and $B = \begin{bmatrix} 2 & 3 \\ 2 & 3 \\ -3 & 0 \end{bmatrix}$, find AB. [2 marks]

9. If $A = \begin{bmatrix} 3 & 6 \\ -3 & 6 \end{bmatrix}$ and $B = \begin{bmatrix} 0 & -5 \\ -1 & -5 \end{bmatrix}$, find AB and BA. Is $AB = BA$? [3 marks]

10. If $A = \begin{bmatrix} 3 & 0 \\ 2 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} -3 & 3 \\ -1 & 0 \end{bmatrix}$, find $(A + B)^2$ and $A^2 + 2AB + B^2$. Are they equal? Why? [5 marks]

11. Find A^T for $A = \begin{bmatrix} -7 \\ -4 \\ -6 \\ 3 \end{bmatrix}$ and state its order. [1 mark]

12. If $A = \begin{bmatrix} 4 & 3 & 6 \\ 3 & 0 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 3 & 3 & 6 \\ -1 & 3 & -5 \end{bmatrix}$, verify that $(4A - B)^T = 4A^T - B^T$. [3 marks]

13. Find x if $\begin{bmatrix} 5 & -6 & -x+6 \\ -6 & -8 & -5 \\ 2 & -5 & 5 \end{bmatrix}$ is symmetric. [1 mark]

14. For $A = \begin{bmatrix} 3 & 3 & -4 \\ -5 & 5 & 6 \\ 6 & -3 & -4 \end{bmatrix}$, find $A - A^T$ and show that it is skew-symmetric. [2 marks]

15. Write $\begin{bmatrix} 3 & 4 & 2 \\ 0 & 3 & 7 \\ -3 & 5 & 3 \end{bmatrix}$ as $P + Q$ where P is symmetric and Q is skew-symmetric. [5 marks]

16. Find the positive number k for which $k\begin{bmatrix} -2 & 1 & 2 \\ -2 & -2 & -1 \\ -1 & 2 & -2 \end{bmatrix}$ is orthogonal, and check one row and two columns of your answer. [5 marks]

17. Show that $A = \begin{bmatrix} -3 & 2 \\ -4 & 3 \end{bmatrix}$ is involutory. [2 marks]

18. Find the transposed conjugate A^θ of $A = \begin{bmatrix} 7-5i & -5-6i \\ 2-3i & 4i-5 \end{bmatrix}$. [1 mark]

19. Show that $A = \begin{bmatrix} 4 & -2 \\ -2 & 4 \end{bmatrix}$ satisfies $A^2 - (a+d)A + (ad-bc)I = O$, writing the equation with numbers. [3 marks]

20. Write the equation $A^3 - (\operatorname{tr} A)A^2 + (\ldots)A - |A|I = O$ satisfied by $A = \begin{bmatrix} 1 & -1 & -2 \\ -3 & -3 & 3 \\ -1 & 3 & -1 \end{bmatrix}$, finding every coefficient. [5 marks]

Key sheet

Q	Answer
1	8 elements; $a_{41} + a_{22} = -9$
2	$A = \begin{bmatrix} 3 & 7 & 11 \\ 2 & 6 & 10 \end{bmatrix}$
3	A scalar matrix
4	6
5	$x = 7, y = 4$
6	$X = \begin{bmatrix} 0 & 3 & 1 \\ -1 & -2 & -4 \end{bmatrix}$
7	$X = \begin{bmatrix} 0 & 5 \\ 0 & 4 \end{bmatrix}, Y = \begin{bmatrix} -1 & 1 \\ 0 & -5 \end{bmatrix}$
8	$AB = \begin{bmatrix} 0 & 18 \end{bmatrix}$
9	$AB = \begin{bmatrix} -6 & -45 \\ -6 & -15 \end{bmatrix}, BA = \begin{bmatrix} 15 & -30 \\ 12 & -36 \end{bmatrix}; AB \neq BA$
10	$(A+B)^2 = \begin{bmatrix} 3 & 3 \\ 1 & 4 \end{bmatrix}, A^2 + 2AB + B^2 = \begin{bmatrix} -3 & 9 \\ -3 & 10 \end{bmatrix}$; not equal since $AB \neq BA$
11	$A^T = \begin{bmatrix} -7 & -4 & -6 & 3 \end{bmatrix}$, order 1×4
12	Both equal $\begin{bmatrix} 13 & 13 \\ 9 & -3 \\ 18 & 9 \end{bmatrix}$
13	$x = 4$
14	$A - A^T = \begin{bmatrix} 0 & 8 & -10 \\ -8 & 0 & 9 \\ 10 & -9 & 0 \end{bmatrix}$
15	$\begin{bmatrix} 3 & 2 & -\frac{1}{2} \\ 2 & 3 & 6 \\ -\frac{1}{2} & 6 & 3 \end{bmatrix} + \begin{bmatrix} 0 & 2 & \frac{5}{2} \\ -2 & 0 & 1 \\ -\frac{5}{2} & -1 & 0 \end{bmatrix}$
16	$k = \frac{1}{3}$
17	$A^2 = I$
18	$A^\theta = \begin{bmatrix} 7+5i & 2+3i \\ 6i-5 & -5-4i \end{bmatrix}$
19	$A^2 - 8A + 12I = O$
20	$A^3 + 3A^2 - 15A - 24I = O$

Solutions

1. For $A = \begin{bmatrix} 0 & 1 \\ -8 & 0 \\ 6 & -5 \\ -9 & 8 \end{bmatrix}$, find the number of elements of A and $a_{41} + a_{22}$. [1]

Order 4×2, so $4 \times 2 = 8$ elements.

$a_{41} + a_{22} = -9 + 0 = -9$

Ans: 8 **elements;** $a_{41} + a_{22} = -9$

---

2. Write the matrix $A = [a_{ij}]_{2\times 3}$ whose elements are $a_{ij} = |i - 4j|$. [2]

$a_{ij} = |i - 4j|$

$a_{11} = 3$, $a_{12} = 7$, $a_{13} = 11$

$a_{21} = 2$, $a_{22} = 6$, $a_{23} = 10$

$A = \begin{bmatrix} 3 & 7 & 11 \\ 2 & 6 & 10 \end{bmatrix}$

Ans: $A = \begin{bmatrix} 3 & 7 & 11 \\ 2 & 6 & 10 \end{bmatrix}$

---

3. Name the type of the square matrix $\begin{bmatrix} -5 & 0 & 0 \\ 0 & -5 & 0 \\ 0 & 0 & -5 \end{bmatrix}$ (the most special type that fits). [1]

Here it is diagonal and all its diagonal elements are equal to -5.

So it is a scalar matrix.

Ans: A scalar matrix

---

4. If $\operatorname{tr}(A) = 1$ and $\operatorname{tr}(B) = 1$, find $\operatorname{tr}(4A + 2B)$. [1]

$\operatorname{tr}(4A + 2B) = 4\operatorname{tr}(A) + 2\operatorname{tr}(B)$

$= 4(1) + 2(1) = 6$

Ans: 6

---

5. If $\begin{bmatrix} 3x - y & 6 \\ 9 & x - y \end{bmatrix} = \begin{bmatrix} 17 & 6 \\ 9 & 3 \end{bmatrix}$, find the values of x and y. [2]

Equate corresponding elements:

$3x - y = 17, \quad x - y = 3$

Solving, $x = 7$, $y = 4$.

Ans: $x = 7$, $y = 4$

---

6. Find the matrix X if $2X + \begin{bmatrix} 3 & -4 & -6 \\ 5 & 7 & 3 \end{bmatrix} = \begin{bmatrix} 3 & 2 & -4 \\ 3 & 3 & -5 \end{bmatrix}$. [2]

$2X = \begin{bmatrix} 3 & 2 & -4 \\ 3 & 3 & -5 \end{bmatrix} - \begin{bmatrix} 3 & -4 & -6 \\ 5 & 7 & 3 \end{bmatrix} = \begin{bmatrix} 0 & 6 & 2 \\ -2 & -4 & -8 \end{bmatrix}$

$X = \frac{1}{2}\begin{bmatrix} 0 & 6 & 2 \\ -2 & -4 & -8 \end{bmatrix} = \begin{bmatrix} 0 & 3 & 1 \\ -1 & -2 & -4 \end{bmatrix}$

Ans: $X = \begin{bmatrix} 0 & 3 & 1 \\ -1 & -2 & -4 \end{bmatrix}$

---

7. Find X and Y if $2X + Y = \begin{bmatrix} -1 & 11 \\ 0 & 3 \end{bmatrix}$ and $X + 2Y = \begin{bmatrix} -2 & 7 \\ 0 & -6 \end{bmatrix}$. [3]

Multiply the first by 2 and subtract the second: $3X = 2\begin{bmatrix} -1 & 11 \\ 0 & 3 \end{bmatrix} - \begin{bmatrix} -2 & 7 \\ 0 & -6 \end{bmatrix} = \begin{bmatrix} 0 & 15 \\ 0 & 12 \end{bmatrix}$, so $X = \begin{bmatrix} 0 & 5 \\ 0 & 4 \end{bmatrix}$.

Then $Y = \begin{bmatrix} -1 & 11 \\ 0 & 3 \end{bmatrix} - 2X = \begin{bmatrix} -1 & 1 \\ 0 & -5 \end{bmatrix}$.

Ans: $X = \begin{bmatrix} 0 & 5 \\ 0 & 4 \end{bmatrix}$, $Y = \begin{bmatrix} -1 & 1 \\ 0 & -5 \end{bmatrix}$

---

8. If $A = \begin{bmatrix} 2 & 4 & 4 \end{bmatrix}$ and $B = \begin{bmatrix} 2 & 3 \\ 2 & 3 \\ -3 & 0 \end{bmatrix}$, find AB. [2]

A is 1×3 and B is 3×2, so AB is 1×2.

$c_{11} = (2)(2) + (4)(2) + (4)(-3) = 0$

$c_{12} = (2)(3) + (4)(3) + (4)(0) = 18$

$AB = \begin{bmatrix} 0 & 18 \end{bmatrix}$

Ans: $AB = \begin{bmatrix} 0 & 18 \end{bmatrix}$

---

9. If $A = \begin{bmatrix} 3 & 6 \\ -3 & 6 \end{bmatrix}$ and $B = \begin{bmatrix} 0 & -5 \\ -1 & -5 \end{bmatrix}$, find AB and BA. Is $AB = BA$? [3]

$AB = \begin{bmatrix} -6 & -45 \\ -6 & -15 \end{bmatrix}$

$BA = \begin{bmatrix} 15 & -30 \\ 12 & -36 \end{bmatrix}$

The products differ, so $AB \neq BA$: matrix multiplication is not commutative.

Ans: $AB = \begin{bmatrix} -6 & -45 \\ -6 & -15 \end{bmatrix}$, $BA = \begin{bmatrix} 15 & -30 \\ 12 & -36 \end{bmatrix}$; $AB \neq BA$

---

10. If $A = \begin{bmatrix} 3 & 0 \\ 2 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} -3 & 3 \\ -1 & 0 \end{bmatrix}$, find $(A+B)^2$ and $A^2 + 2AB + B^2$. Are they equal? Why? [5]

$A + B = \begin{bmatrix} 0 & 3 \\ 1 & 1 \end{bmatrix}$, $(A+B)^2 = \begin{bmatrix} 3 & 3 \\ 1 & 4 \end{bmatrix}$

$A^2 = \begin{bmatrix} 9 & 0 \\ 8 & 1 \end{bmatrix}$, $B^2 = \begin{bmatrix} 6 & -9 \\ 3 & -3 \end{bmatrix}$

$AB = \begin{bmatrix} -9 & 9 \\ -7 & 6 \end{bmatrix}$, $BA = \begin{bmatrix} -3 & 3 \\ -3 & 0 \end{bmatrix}$

$A^2 + 2AB + B^2 = \begin{bmatrix} -3 & 9 \\ -3 & 10 \end{bmatrix}$

They differ because $(A+B)^2 = A^2 + AB + BA + B^2$ and here $AB \neq BA$.

Ans: $(A+B)^2 = \begin{bmatrix} 3 & 3 \\ 1 & 4 \end{bmatrix}$, $A^2 + 2AB + B^2 = \begin{bmatrix} -3 & 9 \\ -3 & 10 \end{bmatrix}$; **not equal since** $AB \neq BA$

---

11. Find A^T for $A = \begin{bmatrix} -7 \\ -4 \\ -6 \\ 3 \end{bmatrix}$ and state its order. [1]

Rows of A become columns of A^T.

$A^T = \begin{bmatrix} -7 & -4 & -6 & 3 \end{bmatrix}$

Order 1×4.

Ans: $A^T = \begin{bmatrix} -7 & -4 & -6 & 3 \end{bmatrix}$, **order** 1×4

---

12. If $A = \begin{bmatrix} 4 & 3 & 6 \\ 3 & 0 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 3 & 3 & 6 \\ -1 & 3 & -5 \end{bmatrix}$, verify that $(4A - B)^T = 4A^T - B^T$. [3]

$4A - B = \begin{bmatrix} 13 & 9 & 18 \\ 13 & -3 & 9 \end{bmatrix}$

$(4A - B)^T = \begin{bmatrix} 13 & 13 \\ 9 & -3 \\ 18 & 9 \end{bmatrix}$

$4A^T - B^T = \begin{bmatrix} 16 & 12 \\ 12 & 0 \\ 24 & 4 \end{bmatrix} - \begin{bmatrix} 3 & -1 \\ 3 & 3 \\ 6 & -5 \end{bmatrix} = \begin{bmatrix} 13 & 13 \\ 9 & -3 \\ 18 & 9 \end{bmatrix}$

The two are equal.

Ans: Both equal $\begin{bmatrix} 13 & 13 \\ 9 & -3 \\ 18 & 9 \end{bmatrix}$

---

13. Find x if $\begin{bmatrix} 5 & -6 & -x+6 \\ -6 & -8 & -5 \\ 2 & -5 & 5 \end{bmatrix}$ is symmetric. [1]

Symmetric means $a_{ij} = a_{ji}$.

$a_{13} = a_{31} \Rightarrow -x + 6 = 2$

$x = 4$

Ans: $x = 4$

---

14. For $A = \begin{bmatrix} 3 & 3 & -4 \\ -5 & 5 & 6 \\ 6 & -3 & -4 \end{bmatrix}$, find $A - A^T$ and show that it is skew-symmetric. [2]

$A^T = \begin{bmatrix} 3 & -5 & 6 \\ 3 & 5 & -3 \\ -4 & 6 & -4 \end{bmatrix}$

$A - A^T = \begin{bmatrix} 0 & 8 & -10 \\ -8 & 0 & 9 \\ 10 & -9 & 0 \end{bmatrix}$

Its diagonal is zero and $a_{ij} = -a_{ji}$, so it is skew-symmetric.

Ans: $A - A^T = \begin{bmatrix} 0 & 8 & -10 \\ -8 & 0 & 9 \\ 10 & -9 & 0 \end{bmatrix}$

---

15. Write $\begin{bmatrix} 3 & 4 & 2 \\ 0 & 3 & 7 \\ -3 & 5 & 3 \end{bmatrix}$ as $P + Q$ where P is symmetric and Q is skew-symmetric. [5]

$A^T = \begin{bmatrix} 3 & 0 & -3 \\ 4 & 3 & 5 \\ 2 & 7 & 3 \end{bmatrix}$

$A + A^T = \begin{bmatrix} 6 & 4 & -1 \\ 4 & 6 & 12 \\ -1 & 12 & 6 \end{bmatrix}$, $P = \frac{1}{2}(A + A^T) = \begin{bmatrix} 3 & 2 & -\frac{1}{2} \\ 2 & 3 & 6 \\ -\frac{1}{2} & 6 & 3 \end{bmatrix}$

$A - A^T = \begin{bmatrix} 0 & 4 & 5 \\ -4 & 0 & 2 \\ -5 & -2 & 0 \end{bmatrix}$, $Q = \frac{1}{2}(A - A^T) = \begin{bmatrix} 0 & 2 & \frac{5}{2} \\ -2 & 0 & 1 \\ -\frac{5}{2} & -1 & 0 \end{bmatrix}$

$P^T = P$ and $Q^T = -Q$, and $P + Q = A$.

$A = \begin{bmatrix} 3 & 2 & -\frac{1}{2} \\ 2 & 3 & 6 \\ -\frac{1}{2} & 6 & 3 \end{bmatrix} + \begin{bmatrix} 0 & 2 & \frac{5}{2} \\ -2 & 0 & 1 \\ -\frac{5}{2} & -1 & 0 \end{bmatrix}$

Ans: $\begin{bmatrix} 3 & 2 & -\frac{1}{2} \\ 2 & 3 & 6 \\ -\frac{1}{2} & 6 & 3 \end{bmatrix} + \begin{bmatrix} 0 & 2 & \frac{5}{2} \\ -2 & 0 & 1 \\ -\frac{5}{2} & -1 & 0 \end{bmatrix}$

---

16. Find the positive number k for which $k \begin{bmatrix} -2 & 1 & 2 \\ -2 & -2 & -1 \\ -1 & 2 & -2 \end{bmatrix}$ is orthogonal, and check one row and two columns of your answer. [5]

Let $A = kM$. Then $AA^T = k^2 MM^T$.

$MM^T = \begin{bmatrix} 9 & 0 & 0 \\ 0 & 9 & 0 \\ 0 & 0 & 9 \end{bmatrix} = 9I$

$AA^T = I \Rightarrow 9k^2 = 1 \Rightarrow k = \frac{1}{3}$

Row 1: $\frac{1}{9}((-2)^2 + 1^2 + 2^2) = 1$.

Columns 1, 2: $\frac{1}{9}((-2)(1) + (-2)(-2) + (-1)(2)) = 0$.

Ans: $k = \frac{1}{3}$

---

17. Show that $A = \begin{bmatrix} -3 & 2 \\ -4 & 3 \end{bmatrix}$ is involutory. [2]

$A^2 = \begin{bmatrix} -3 & 2 \\ -4 & 3 \end{bmatrix} \begin{bmatrix} -3 & 2 \\ -4 & 3 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

$A^2 = I$, so A is involutory.

Ans: $A^2 = I$

---

18. Find the transposed conjugate A^θ of $A = \begin{bmatrix} 7 - 5i & -5 - 6i \\ 2 - 3i & 4i - 5 \end{bmatrix}$. [1]

$\bar{A} = \begin{bmatrix} 7 + 5i & 6i - 5 \\ 2 + 3i & -5 - 4i \end{bmatrix}$

$A^\theta = (\bar{A})^T = \begin{bmatrix} 7 + 5i & 2 + 3i \\ 6i - 5 & -5 - 4i \end{bmatrix}$

Ans: $A^\theta = \begin{bmatrix} 7 + 5i & 2 + 3i \\ 6i - 5 & -5 - 4i \end{bmatrix}$

---

19. Show that $A = \begin{bmatrix} 4 & -2 \\ -2 & 4 \end{bmatrix}$ satisfies $A^2 - (a+d)A + (ad-bc)I = O$, writing the equation with numbers. [3]

$a + d = 8, \quad ad - bc = 12$

$A^2 = \begin{bmatrix} 20 & -16 \\ -16 & 20 \end{bmatrix}$

$A^2 - (8)A + (12)I = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$

$A^2 - 8A + 12I = O$

Ans: $A^2 - 8A + 12I = O$

---

20. Write the equation $A^3 - (\operatorname{tr} A)A^2 + (\ldots)A - |A|I = O$ satisfied by $A = \begin{bmatrix} 1 & -1 & -2 \\ -3 & -3 & 3 \\ -1 & 3 & -1 \end{bmatrix}$, finding every coefficient. [5]

$\operatorname{tr} A = 1 - 3 - 1 = -3$

$\begin{vmatrix} -3 & 3 \\ 3 & -1 \end{vmatrix} = -6, \quad \begin{vmatrix} 1 & -2 \\ -1 & -1 \end{vmatrix} = -3, \quad \begin{vmatrix} 1 & -1 \\ -3 & -3 \end{vmatrix} = -6$

Sum of the principal minors $= -15$.

$|A| = 24$

So $A^3 + 3A^2 - 15A - 24I = O$.

Ans: $A^3 + 3A^2 - 15A - 24I = O$