

Polynomials

MCQ Test, Set 04

Time: 30 minutes

Maximum marks: 25

Name: _____ Class: _____ Date: _____

- There are 25 questions. Each has exactly one correct answer: choose A, B, C or D.
- Each correct answer carries 1 mark. There is no negative marking.
- The key sheet and worked solutions are at the end of this test.

1. Which one of these is a monomial?

- A.** $-6x^4y^2$ **B.** $\frac{5}{x^4}$ **C.** $\frac{7x}{y}$ **D.** $-2\sqrt{xy}^2$

2. The degree of $-3y^3 - 5x^5y^2 - 8x^2 + 5$ is

- A.** 4 **B.** 3 **C.** 7 **D.** 5

3. When $p(x) = 9x - 9 + x^4$ is written in standard form $a_nx^n + \dots + a_1x + a_0$, the value of a_n is

- A.** 9 **B.** 1 **C.** -9 **D.** 4

4. Which of the following is a quadratic polynomial?

- A.** $8x - 7x^2 - 4x^3 - 3$ **B.** $-6 + 2x$ **C.** $-9x^3 + 3x^4 - 8$ **D.** $x - 8x^2 + 7$

5. Among the following, the one that is not a polynomial is

- A.** $-4x^{-3} - 9x^2 + 2$ **B.** $-3\sqrt{5}x^4 - 6x^2 - 2\sqrt{5}x^3$
C. $4\sqrt{5}x^4 + 4\sqrt{3}x + 8x^3$ **D.** $2x^4 + \frac{7}{4}x^2$

6. If $(p+5)x^2 + (q+2p)x + (r+2q)$ is a zero polynomial, then

- A.** $p = -5, q = 10, r = -20$ **B.** $p = -5, q = -10, r = 20$ **C.** $p = -5, q = 10, r = 20$ **D.** $p = 5, q = -10, r = 20$

7. Pick the quadratic equation.

- A.** $-6x - 1 = 0$ **B.** $x^2 - 4\sqrt{x} + 7 = 0$
C. $5\sqrt{2}x^2 - x + 4\sqrt{2} = 0$ **D.** $x^2 + \frac{2}{x^2} = 7$

8. If $p(x) = x^3 + 5x^2 + 7x - 8$, then $p(-1) =$

- A.** -11 **B.** 5 **C.** 3 **D.** -21

9. If 1 is a zero of $p(x) = x^3 + 3x^2 - ax - 5$, then $a =$

- A.** 3 **B.** 1 **C.** 0 **D.** -1

10. The roots of $5x^2 + 19x + 12 = 0$ are

- A.** $-3, -\frac{4}{5}$ **B.** $-3, -4$
C. $-\frac{1}{3}, -\frac{5}{4}$ **D.** $3, \frac{4}{5}$

11. The roots of $x^2 - 10x + 9 = 0$ are
A. 1, -9 **B.** -1, 9 **C.** 1, 9 **D.** -1, -9
-
12. The roots of $x^2 - 10x - 5 = 0$ are
A. $-5 \pm \sqrt{30}$ **B.** $5 \pm \sqrt{30}$ **C.** $10 \pm 2\sqrt{30}$ **D.** $5 \pm 2\sqrt{5}$
-
13. The remainder when $x^4 + 4x^3 + x^2 - 10x - 8$ is divided by $x^2 - x - 1$ is
A. $2x + 1$ **B.** $-2x - 1$ **C.** $-2x + 1$ **D.** $2x - 1$
-
14. The remainder when $16x^3 - 8x^2 + 4x + 5$ is divided by $2x + 3$ is
A. -511 **B.** 73 **C.** -73 **D.** 47
-
15. If $x + 2$ is a factor of $x^3 + 5x^2 + ax + 4$, then $a =$
A. 8 **B.** -16 **C.** 6 **D.** -8
-
16. $3\sqrt{2}x^2 + 7x - 3\sqrt{2} =$
A. $(\sqrt{2}x - 3)(3x - \sqrt{2})$ **B.** $(\sqrt{2}x + 3)(3x + \sqrt{2})$
C. $(\sqrt{2}x - 3)(3x + \sqrt{2})$ **D.** $(\sqrt{2}x + 3)(3x - \sqrt{2})$
-
17. What must be subtracted from $-3x^3 + 6x^2 + 8x - 1$ so that it is exactly divisible by $x^2 - 1$?
A. $-3x + 6$ **B.** $5x + 5$ **C.** $5x - 5$ **D.** $-5x - 5$
-
18. If $x - 4$ and $x + 2$ are factors of $x^3 + ax^2 + bx + 8$, then
A. $a = 3, b = 6$ **B.** $a = 3, b = -6$ **C.** $a = -3, b = 6$ **D.** $a = -3, b = -6$
-
19. $x^2 + 4x - 7 =$
A. $(x + 2 - \sqrt{11})(x + 2 + \sqrt{11})$ **B.** $(x - 2 - \sqrt{11})(x - 2 + \sqrt{11})$
C. $(x + 2 - 2\sqrt{11})(x + 2 + 2\sqrt{11})$ **D.** $(x + 4 - \sqrt{11})(x + 4 + \sqrt{11})$
-
20. Given that $x + 5$ is a factor, $x^3 - 19x + 30 =$
A. $(x + 5)(x^2 - 5x - 6)$ **B.** $(x + 5)(x^2 + 5x - 6)$ **C.** $(x + 5)(x^2 - 5x + 6)$ **D.** $(x + 5)(x^2 + 5x + 6)$
-
21. $x^3 - 3x^2 - 10x + 24 =$
A. $(x - 4)(x - 3)(x - 2)$ **B.** $(x - 4)(x + 3)(x - 2)$ **C.** $(x + 4)(x - 3)(x + 2)$ **D.** $(x + 4)(x + 3)(x - 2)$
-
22. Using synthetic division, the remainder when $x^3 + 3x^2 + 5$ is divided by $x - 1$ is
A. 4 **B.** 9 **C.** 7 **D.** -9
-
23. $x^4 - 81 = (x - 3) \times$
A. $x^3 + 3x^2 + 3x + 3$ **B.** $x^3 + 3x^2 + 6x + 9$ **C.** $x^3 - 3x^2 + 9x - 27$ **D.** $x^3 + 3x^2 + 9x + 27$
-

Key sheet

1. A	2. C	3. B	4. D	5. A
6. A	7. C	8. A	9. D	10. A
11. C	12. B	13. D	14. C	15. A
16. D	17. B	18. D	19. A	20. C
21. B	22. B	23. D	24. C	25. B

Solutions

1. Which one of these is a monomial?

$-6x^4y^2$: the powers of x and y are 4 and 2, non-negative integers, so it is a monomial.
Each of the others has a negative or fractional power.

Ans: (A) $-6x^4y^2$

2. The degree of $-3y^3 - 5x^5y^2 - 8x^2 + 5$ is

Degrees of the terms: 3, 7, 2, 0

The greatest is 7.

Ans: (C) 7

3. When $p(x) = 9x - 9 + x^4$ is written in standard form $a_nx^n + \dots + a_1x + a_0$, the value of a_n is

Standard form: $x^4 + 9x - 9$

$n = 4$ and a_n is the coefficient of x^4 : $a_n = 1$

Ans: (B) 1

4. Which of the following is a quadratic polynomial?

$x - 8x^2 + 7$ has highest power 2, so it is a quadratic polynomial.

Ans: (D) $x - 8x^2 + 7$

5. Among the following, the one that is not a polynomial is

In $-4x^{-3} - 9x^2 + 2$, the power -3 of x is negative.

So it is not a polynomial. (A coefficient like $\sqrt{2}$ or $\frac{2}{3}$ is allowed.)

Ans: (A) $-4x^{-3} - 9x^2 + 2$

6. If $(p + 5)x^2 + (q + 2p)x + (r + 2q)$ is a zero polynomial, then

$$p + 5 = 0 \Rightarrow p = -5$$

$$q + 2p = 0 \Rightarrow q - 10 = 0 \Rightarrow q = 10$$

$$r + 2q = 0 \Rightarrow r + 20 = 0 \Rightarrow r = -20$$

Ans: (A) $p = -5, q = 10, r = -20$

7. Pick the quadratic equation.

$$5\sqrt{2}x^2 - x + 4\sqrt{2} = 0: \text{ it has the form } ax^2 + bx + c = 0 \text{ with } a \neq 0.$$

The others have a negative or fractional power of x , or a degree other than 2.

Ans: (C) $5\sqrt{2}x^2 - x + 4\sqrt{2} = 0$

8. If $p(x) = x^3 + 5x^2 + 7x - 8$, then $p(-1) =$

$$p(-1) = (-1)^3 + 5(-1)^2 + 7(-1) - 8$$

$$= -1 + 5 - 7 - 8$$

$$= -11$$

Ans: (A) -11

9. If 1 is a zero of $p(x) = x^3 + 3x^2 - ax - 5$, then $a =$

$$p(1) = (1)^3 + 3(1)^2 - a(1) - 5$$

$$= 1 + 3 - a - 5$$

$$= -a - 1$$

$$-a - 1 = 0 \Rightarrow a = -1$$

Ans: (D) -1

10. The roots of $5x^2 + 19x + 12 = 0$ are

$$5x^2 + 19x + 12 = 5x^2 + 4x + 15x + 12$$

$$= x(5x + 4) + 3(5x + 4)$$

$$= (5x + 4)(x + 3)$$

$$\Rightarrow x = -3, -\frac{4}{5}$$

Ans: (A) $-3, -\frac{4}{5}$

11. The roots of $x^2 - 10x + 9 = 0$ are

$$x^2 - 10x + 9 = x^2 - x - 9x + 9$$

$$= x(x - 1) - 9(x - 1)$$

$$= (x - 1)(x - 9)$$

$$\Rightarrow x = 1, 9$$

Ans: (C) $1, 9$

12. The roots of $x^2 - 10x - 5 = 0$ are

$$a = 1, b = -10, c = -5$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{10 \pm \sqrt{100 - 4(1)(-5)}}{2(1)}$$

$$= 5 + \sqrt{30}, 5 - \sqrt{30}$$

Ans: (B) $5 \pm \sqrt{30}$

13. The remainder when $x^4 + 4x^3 + x^2 - 10x - 8$ is divided by $x^2 - x - 1$ is

$$x^4 \div x^2 = x^2; x^2 \times (x^2 - x - 1) = x^4 - x^3 - x^2; \text{ subtract: } 5x^3 + 2x^2 - 10x - 8$$

$$5x^3 \div x^2 = 5x; 5x \times (x^2 - x - 1) = 5x^3 - 5x^2 - 5x; \text{ subtract: } 7x^2 - 5x - 8$$

$$7x^2 \div x^2 = 7; 7 \times (x^2 - x - 1) = 7x^2 - 7x - 7; \text{ subtract: } 2x - 1$$

$$\text{Remainder} = 2x - 1$$

Ans: (D) $2x - 1$

14. The remainder when $16x^3 - 8x^2 + 4x + 5$ is divided by $2x + 3$ is

$$2x + 3 = 0 \Rightarrow x = -\frac{3}{2}; \text{ remainder} = f\left(-\frac{3}{2}\right)$$

$$= -73$$

Ans: (C) -73

15. If $x + 2$ is a factor of $x^3 + 5x^2 + ax + 4$, then $a =$

$$\begin{aligned} f(-2) &= (-2)^3 + 5(-2)^2 + a(-2) + 4 \\ &= -8 + 20 - 2a + 4 \\ &= -2a + 16 \\ -2a + 16 &= 0 \Rightarrow a = 8 \end{aligned}$$

Ans: (A) 8

16. $3\sqrt{2}x^2 + 7x - 3\sqrt{2} =$

$$\begin{aligned} 3\sqrt{2}x^2 + 7x - 3\sqrt{2} &= 3\sqrt{2}x^2 - 2x + 9x - 3\sqrt{2} \\ &= \sqrt{2}x(3x - \sqrt{2}) + 3(3x - \sqrt{2}) = (\sqrt{2}x + 3)(3x - \sqrt{2}) \end{aligned}$$

Ans: (D) $(\sqrt{2}x + 3)(3x - \sqrt{2})$

17. What must be subtracted from $-3x^3 + 6x^2 + 8x - 1$ so that it is exactly divisible by $x^2 - 1$?

$$\begin{aligned} -3x^3 \div x^2 &= -3x; (-3x) \times (x^2 - 1) = -3x^3 + 3x; \text{ subtract: } 6x^2 + 5x - 1 \\ 6x^2 \div x^2 &= 6; 6 \times (x^2 - 1) = 6x^2 - 6; \text{ subtract: } 5x + 5 \\ \text{Remainder} &= 5x + 5, \text{ so } 5x + 5 \text{ must be subtracted.} \end{aligned}$$

Ans: (B) $5x + 5$

18. If $x - 4$ and $x + 2$ are factors of $x^3 + ax^2 + bx + 8$, then

$$\begin{aligned} f(4) &= 0 \Rightarrow 4a + b = -18 \dots (1) \\ f(-2) &= 0 \Rightarrow 2a - b = 0 \dots (2) \\ (1) + (2): 6a &= -18 \\ \Rightarrow a &= -3 \\ \text{Put } a = -3 \text{ in (1): } -12 + b &= -18 \Rightarrow b = -6 \end{aligned}$$

Ans: (D) $a = -3, b = -6$

19. $x^2 + 4x - 7 =$

$$\begin{aligned} x &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-4 \pm \sqrt{16 - 4(1)(-7)}}{2(1)} \\ &= \frac{-4 \pm \sqrt{44}}{2} \\ &= \frac{-4 \pm 2\sqrt{11}}{2} \\ &= -2 \pm \sqrt{11} \\ &= -2 + \sqrt{11}, -2 - \sqrt{11} \\ \text{So } x^2 + 4x - 7 &= (x - \alpha)(x - \beta) = (x + 2 - \sqrt{11})(x + 2 + \sqrt{11}) \end{aligned}$$

Ans: (A) $(x + 2 - \sqrt{11})(x + 2 + \sqrt{11})$

20. Given that $x + 5$ is a factor, $x^3 - 19x + 30 =$

$$\begin{aligned} x^3 \div x &= x^2; x^2 \times (x + 5) = x^3 + 5x^2; \text{ subtract: } -5x^2 - 19x + 30 \\ -5x^2 \div x &= -5x; (-5x) \times (x + 5) = -5x^2 - 25x; \text{ subtract: } 6x + 30 \\ 6x \div x &= 6; 6 \times (x + 5) = 6x + 30; \text{ subtract: } 0 \\ \text{Quotient} &= x^2 - 5x + 6 \end{aligned}$$

Ans: (C) $(x + 5)(x^2 - 5x + 6)$

21. $x^3 - 3x^2 - 10x + 24 =$

$$f(2) = 8 - 12 - 20 + 24 = 0$$

Dividing and splitting the quotient gives $f(x) = (x - 4)(x + 3)(x - 2)$.

Ans: (B) $(x - 4)(x + 3)(x - 2)$

22. Using synthetic division, the remainder when $x^3 + 3x^2 + 5$ is divided by $x - 1$ is

Second row: 0 1 4 4

Bottom row: 1 4 4 | 9

Remainder = 9

Ans: (B) 9

23. $x^4 - 81 = (x - 3) \times$

$$x^4 - 81 = (x - 3)(x^3 + 3x^2 + 9x + 27)$$

Signs are all +; the powers of 3 go up as the powers of x go down.

Ans: (D) $x^3 + 3x^2 + 9x + 27$

24. When $8a^3 + 6a^2x + \frac{3}{2}ax^2 + \frac{1}{8}x^3$ is divided by $2a + \frac{1}{2}x$, the quotient is

$$8a^3 \div 2a = 4a^2; 4a^2 \times (2a + \frac{1}{2}x) = 8a^3 + 2a^2x; \text{ subtract: } 4a^2x + \frac{3}{2}ax^2 + \frac{1}{8}x^3$$

$$4a^2x \div 2a = 2ax; 2ax \times (2a + \frac{1}{2}x) = 4a^2x + ax^2; \text{ subtract: } \frac{1}{2}ax^2 + \frac{1}{8}x^3$$

$$\frac{1}{2}ax^2 \div 2a = \frac{1}{4}x^2; \frac{1}{4}x^2 \times (2a + \frac{1}{2}x) = \frac{1}{2}ax^2 + \frac{1}{8}x^3; \text{ subtract: } 0$$

Ans: (C) $4a^2 + 2ax + \frac{1}{4}x^2$

25. The quadratic polynomial which leaves the remainders -7 , -10 and -11 when divided by $x - 1$, $x - 2$ and $x - 3$ respectively is

Check each remainder with the remainder theorem: $f(1) = -7$, $f(2) = -10$, $f(3) = -11$.

Only $x^2 - 6x - 2$ gives all three.

Ans: (B) $x^2 - 6x - 2$
