

Polynomials

MCQ Test, Set 05

Time: 30 minutes

Maximum marks: 25

Name: _____ Class: _____ Date: _____

- There are 25 questions. Each has exactly one correct answer: choose A, B, C or D.
- Each correct answer carries 1 mark. There is no negative marking.
- The key sheet and worked solutions are at the end of this test.

1. Which one of these is NOT a monomial?

- A.** $-\frac{7}{5}x^5y^3$ **B.** $6x^{\frac{1}{3}}y^3$ **C.** $2x^5y^4$ **D.** $-8x^2y^2$

2. The degree of the monomial $-2x^2y^3z^3$ is

- A.** 8 **B.** 3 **C.** 18 **D.** 9

3. The coefficient of x^2 in $9x^2 + 5x^3 + 2 - 9x^4$ is

- A.** 2 **B.** -9 **C.** 9 **D.** 5

4. $-7x + 6 + 9x^4 - x^3 - 7x^2$ is a

- A.** Quadratic polynomial **B.** Bi-quadratic polynomial
C. Linear polynomial **D.** Cubic polynomial

5. Which one of these expressions is a polynomial?

- A.** $8x^{\frac{1}{3}} - 2x^3 - \frac{5}{3}x$ **B.** $-2\sqrt{3}x^2 + \frac{5}{3}x + 7\sqrt{x}$
C. $2\sqrt{x} + 6$ **D.** $-3\sqrt{5}x^5 + 5$

6. If $5x^{n-9} + 5x^{n-10} - 6x^{n-11}$ is a quadratic polynomial ($n \in N$), then $n =$

- A.** 11 **B.** 12 **C.** 2 **D.** 13

7. The equation $-4x^4 + 7x^2 + 6x + 7 = 0$ is a

- A.** Cubic equation **B.** Bi-quadratic equation
C. Quadratic equation **D.** Linear equation

8. If $p(x) = 4x^2 + 7x + 10$, then $p(-3) =$

- A.** -47 **B.** 67 **C.** 25 **D.** 5

9. If $p(x) = -2x^3 - 5x^2 - 4x - 4$, then $p(-3) =$

- A.** 17 **B.** -115 **C.** -7 **D.** 107

10. The zero of the polynomial $4x + 11$ is

- A.** $-\frac{11}{4}$ **B.** $\frac{4}{11}$ **C.** $-\frac{4}{11}$ **D.** $\frac{11}{4}$

11. The roots of $15x^2 + 13x + 2 = 0$ are
A. $-2, -1$ **B.** $\frac{2}{3}, \frac{1}{5}$
C. $-\frac{2}{3}, -\frac{1}{5}$ **D.** $-\frac{3}{2}, -5$
-
12. The discriminant of $3x^2 + 10x - 11 = 0$ is
A. -32 **B.** 32 **C.** 133 **D.** 232
-
13. A polynomial of degree 6 is divided by a polynomial of degree 4. The degree of the quotient is
A. 4 **B.** 24 **C.** 10 **D.** 2
-
14. If $x^3 + ax^2 - 5x - 6$ leaves the remainder -2 when divided by $x - 1$, then $a =$
A. 0 **B.** 8 **C.** 12 **D.** 10
-
15. If $16x^4 + 4x^3 - ax^2 + 3x - 4$ is divisible by $1 - 2x$, then $a =$
A. 19 **B.** 4 **C.** -4 **D.** -20
-
16. $5x^2 + 37x - 24 =$
A. $(x + 3)(5x - 8)$ **B.** $(5x - 3)(x + 8)$ **C.** $(x - 3)(5x + 8)$ **D.** $(x - 8)(5x + 3)$
-
17. What must be subtracted from $x^3 - 2x^2 + x - 5$ so that it is exactly divisible by $x - 4$?
A. 31 **B.** -105 **C.** 105 **D.** -31
-
18. If $x - 2$ and $x - 3$ are factors of $x^3 + ax^2 + bx + 6$, then
A. $a = 4, b = 1$ **B.** $a = 4, b = -1$ **C.** $a = -4, b = -1$ **D.** $a = -4, b = 1$
-
19. A quadratic polynomial has zeros 3 and $\frac{5}{3}$ and leading coefficient 3. The polynomial is
A. $3x^2 + 14x + 15$ **B.** $3x^2 - 14x + 15$
C. $3x^2 - 4x - 15$ **D.** $x^2 - \frac{14}{3}x + 5$
-
20. Given that $x - 4$ is a factor, $x^3 - 5x^2 + 2x + 8 =$
A. $(x - 4)(x^2 - 9x + 38)$ **B.** $(x - 4)(x^2 - x + 2)$ **C.** $(x - 4)(x^2 + x - 2)$ **D.** $(x - 4)(x^2 - x - 2)$
-
21. $x^3 + 8x^2 + 19x + 12 =$
A. $(x + 3)(x + 4)(x + 1)$ **B.** $(x + 3)(x - 4)(x + 1)$ **C.** $(x - 3)(x - 4)(x - 1)$ **D.** $(x - 3)(x + 4)(x + 1)$
-
22. Using synthetic division, the quotient when $x^3 - 2x + 4$ is divided by $x - 4$ is
A. $x^3 + 4x^2 + 14x + 60$ **B.** $x + 2$
C. $x^2 - 4x + 14$ **D.** $x^2 + 4x + 14$
-
23. $22^{93} - 13^{93}$ is exactly divisible by
A. 13 **B.** 9 **C.** 22 **D.** 35
-

24. When $\frac{1}{8}a^3 - \frac{9}{4}a^2x + \frac{27}{2}ax^2 - 27x^3$ is divided by $\frac{1}{2}a - 3x$, the quotient is

A. $\frac{1}{4}a^2 + 3ax + 9x^2$

B. $\frac{1}{4}a^2 + 9x^2$

C. $\frac{1}{4}a^2 - 3ax + 9x^2$

D. $\frac{1}{4}a^2 - 3ax - 9x^2$

25. $x^2 + bx + c$ leaves the remainders -4 and 5 when divided by $x - 1$ and $x + 2$ respectively. Then

A. $b = 2, c = -3$

B. $b = 2, c = 3$

C. $b = -2, c = -3$

D. $b = -2, c = 3$

Key sheet

1. B	2. A	3. C	4. B	5. D
6. A	7. B	8. C	9. A	10. A
11. C	12. D	13. D	14. B	15. C
16. B	17. A	18. D	19. B	20. D
21. A	22. D	23. B	24. C	25. C

Solutions

1. Which one of these is NOT a monomial?

The power $\frac{1}{3}$ of x is not a whole number.

So $6x^{\frac{1}{3}}y^3$ is not a monomial; the others have only non-negative integer powers.

Ans: (B) $6x^{\frac{1}{3}}y^3$

2. The degree of the monomial $-2x^2y^3z^3$ is

Degree = sum of the exponents = $2 + 3 + 3 = 8$

Ans: (A) 8

3. The coefficient of x^2 in $9x^2 + 5x^3 + 2 - 9x^4$ is

The term with x^2 is $9x^2$, so the coefficient is 9.

Ans: (C) 9

4. $-7x + 6 + 9x^4 - x^3 - 7x^2$ is a

The highest power of x is 4, so it is a bi-quadratic polynomial.

Ans: (B) Bi-quadratic polynomial

5. Which one of these expressions is a polynomial?

In $-3\sqrt{5}x^5 + 5$, all powers of x are whole numbers, so it is a polynomial.

Each other option has a negative or fractional power of x .

Ans: (D) $-3\sqrt{5}x^5 + 5$

6. If $5x^{n-9} + 5x^{n-10} - 6x^{n-11}$ is a quadratic polynomial ($n \in \mathbb{N}$), then $n =$

The degree of a quadratic polynomial is 2.

So the highest power: $n - 9 = 2$

$\Rightarrow n = 11$

Ans: (A) 11

7. The equation $-4x^4 + 7x^2 + 6x + 7 = 0$ is a

The highest power of x is 4, so it is a bi-quadratic equation.

Ans: (B) Bi-quadratic equation

8. If $p(x) = 4x^2 + 7x + 10$, then $p(-3) =$

$p(-3) = 4(-3)^2 + 7(-3) + 10$

$= 36 - 21 + 10$

$= 25$

Ans: (C) 25

9. If $p(x) = -2x^3 - 5x^2 - 4x - 4$, then $p(-3) =$
 $p(-3) = -2(-3)^3 - 5(-3)^2 - 4(-3) - 4$
 $= 54 - 45 + 12 - 4$
 $= 17$

Ans: (A) 17

10. The zero of the polynomial $4x + 11$ is
 $4x + 11 = 0 \Rightarrow 4x = -11 \Rightarrow x = -\frac{11}{4}$

Ans: (A) $-\frac{11}{4}$

11. The roots of $15x^2 + 13x + 2 = 0$ are
 $15x^2 + 13x + 2 = 15x^2 + 3x + 10x + 2$
 $= 3x(5x + 1) + 2(5x + 1)$
 $= (5x + 1)(3x + 2)$
 $\Rightarrow x = -\frac{2}{3}, -\frac{1}{5}$

Ans: (C) $-\frac{2}{3}, -\frac{1}{5}$

12. The discriminant of $3x^2 + 10x - 11 = 0$ is
 $\Delta = b^2 - 4ac = 10^2 - 4(3)(-11) = 232$

Ans: (D) 232

13. A polynomial of degree 6 is divided by a polynomial of degree 4. The degree of the quotient is
 $f(x) = g(x)q(x) + r(x)$: degree of $q(x) = \text{degree of } f(x) - \text{degree of } g(x)$
 $= 6 - 4 = 2$

Ans: (D) 2

14. If $x^3 + ax^2 - 5x - 6$ leaves the remainder -2 when divided by $x - 1$, then $a =$
 $f(1) = (1)^3 + a(1)^2 - 5(1) - 6$
 $= 1 + a - 5 - 6$
 $= a - 10$
 $a - 10 = -2 \Rightarrow a = 8$

Ans: (B) 8

15. If $16x^4 + 4x^3 - ax^2 + 3x - 4$ is divisible by $1 - 2x$, then $a =$
 $1 - 2x = 0 \Rightarrow x = \frac{1}{2}; f\left(\frac{1}{2}\right) = 0$
 $f\left(\frac{1}{2}\right) = 16\left(\frac{1}{2}\right)^4 + 4\left(\frac{1}{2}\right)^3 - a\left(\frac{1}{2}\right)^2 + 3\left(\frac{1}{2}\right) - 4$
 $= 1 + \frac{1}{2} - \frac{1}{4}a + \frac{3}{2} - 4$
 $= -\frac{1}{4}a - 1$
 $-\frac{1}{4}a - 1 = 0 \Rightarrow a = -4$

Ans: (C) -4

16. $5x^2 + 37x - 24 =$

$$\begin{aligned}5x^2 + 37x - 24 &= 5x^2 - 3x + 40x - 24 \\&= x(5x - 3) + 8(5x - 3) \\&= (5x - 3)(x + 8)\end{aligned}$$

Ans: (B) $(5x - 3)(x + 8)$

17. What must be subtracted from $x^3 - 2x^2 + x - 5$ so that it is exactly divisible by $x - 4$?

Remainder = $f(4)$ (remainder theorem)

$$\begin{aligned}f(4) &= (4)^3 - 2(4)^2 + (4) - 5 \\&= 64 - 32 + 4 - 5 \\&= 31\end{aligned}$$

The remainder is 31, so 31 must be subtracted.

Ans: (A) 31

18. If $x - 2$ and $x - 3$ are factors of $x^3 + ax^2 + bx + 6$, then

$$f(2) = 0 \Rightarrow 2a + b = -7 \dots (1)$$

$$f(3) = 0 \Rightarrow 3a + b = -11 \dots (2)$$

$$(1) - (2): -a = 4$$

$$\Rightarrow a = -4$$

$$\text{Put } a = -4 \text{ in (1): } -8 + b = -7 \Rightarrow b = 1$$

Ans: (D) $a = -4, b = 1$

19. A quadratic polynomial has zeros 3 and $\frac{5}{3}$ and leading coefficient 3. The polynomial is

$$\begin{aligned}\text{With zeros } 3, \frac{5}{3} \text{ and leading coefficient } 3: & 3(x - 3)\left(x - \frac{5}{3}\right) \\&= 3x^2 - 14x + 15\end{aligned}$$

Ans: (B) $3x^2 - 14x + 15$

20. Given that $x - 4$ is a factor, $x^3 - 5x^2 + 2x + 8 =$

$$x^3 \div x = x^2; x^2 \times (x - 4) = x^3 - 4x^2; \text{ subtract: } -x^2 + 2x + 8$$

$$-x^2 \div x = -x; (-x) \times (x - 4) = -x^2 + 4x; \text{ subtract: } -2x + 8$$

$$-2x \div x = -2; (-2) \times (x - 4) = -2x + 8; \text{ subtract: } 0$$

$$\text{Quotient} = x^2 - x - 2$$

Ans: (D) $(x - 4)(x^2 - x - 2)$

21. $x^3 + 8x^2 + 19x + 12 =$

$$f(-1) = -1 + 8 - 19 + 12 = 0$$

Dividing and splitting the quotient gives $f(x) = (x + 3)(x + 4)(x + 1)$.

Ans: (A) $(x + 3)(x + 4)(x + 1)$

22. Using synthetic division, the quotient when $x^3 - 2x + 4$ is divided by $x - 4$ is

Second row: 0 4 16 56

Bottom row: 1 4 14 | 60

Quotient = $x^2 + 4x + 14$

Ans: (D) $x^2 + 4x + 14$

23. $22^{93} - 13^{93}$ is exactly divisible by

$n = 93$: $x - a$ is a factor of $x^n - a^n$ (n odd).

So $22 - 13 = 9$ divides it.

Ans: (B) 9

24. When $\frac{1}{8}a^3 - \frac{9}{4}a^2x + \frac{27}{2}ax^2 - 27x^3$ is divided by $\frac{1}{2}a - 3x$, the quotient is

$\frac{1}{8}a^3 \div \frac{1}{2}a = \frac{1}{4}a^2$; $\frac{1}{4}a^2 \times (\frac{1}{2}a - 3x) = \frac{1}{8}a^3 - \frac{3}{4}a^2x$; subtract: $-\frac{3}{2}a^2x + \frac{27}{2}ax^2 - 27x^3$
 $-\frac{3}{2}a^2x \div \frac{1}{2}a = -3ax$; $(-3ax) \times (\frac{1}{2}a - 3x) = -\frac{3}{2}a^2x + 9ax^2$; subtract: $\frac{9}{2}ax^2 - 27x^3$
 $\frac{9}{2}ax^2 \div \frac{1}{2}a = 9x^2$; $9x^2 \times (\frac{1}{2}a - 3x) = \frac{9}{2}ax^2 - 27x^3$; subtract: 0

Ans: (C) $\frac{1}{4}a^2 - 3ax + 9x^2$

25. $x^2 + bx + c$ leaves the remainders -4 and 5 when divided by $x - 1$ and $x + 2$ respectively. Then

$f(1) = 1 + b + c = -4 \Rightarrow b + c = -5 \dots (1)$

$f(-2) = 4 - 2b + c = 5 \Rightarrow 2b - c = -1 \dots (2)$

$(1) + (2)$: $3b = -6$

$\Rightarrow b = -2$

Put $b = -2$ in (1): $-2 + c = -5 \Rightarrow c = -3$

Ans: (C) $b = -2, c = -3$
