

Rationalization (Surds)

MCQ Test, Set 01

Time: 30 minutes

Maximum marks: 25

Name: _____ Class: _____ Date: _____

- There are 25 questions. Each has exactly one correct answer: choose A, B, C or D.
- Each correct answer carries 1 mark. There is no negative marking.
- The key sheet and worked solutions are at the end of this test.

1. Which of the following is a rational number?

- A. $\sqrt[3]{64}$ B. $\sqrt{54}$ C. $\sqrt{12}$ D. $\sqrt[4]{17}$

2. $\sqrt[6]{15}$ and e are both irrational. Which of them is a surd?

- A. e only B. $\sqrt[6]{15}$ only C. Both D. Neither

3. $9\sqrt[3]{n}$ is a mixed surd when n is

- A. 64 B. 729 C. 512 D. 19

4. The conjugate surd of $12 - \sqrt{30}$ is

- A. $-12 - \sqrt{30}$ B. $12 + \sqrt{30}$ C. $-12 + \sqrt{30}$ D. $\frac{1}{12 - \sqrt{30}}$

5. $\sqrt{10} \times \sqrt{10} =$

- A. $2\sqrt{10}$ B. 10 C. $\sqrt{10}$ D. 100

6. $\frac{\sqrt{171}}{\sqrt{19}} =$

- A. $\sqrt{152}$ B. 9 C. 3 D. $3\sqrt{19}$

7. $64^{\frac{1}{6}} =$

- A. 4 B. 2 C. $\frac{64}{6}$ D. 12

8. $\frac{\sqrt[4]{80}}{\sqrt[4]{5}} =$

- A. 2 B. $\sqrt[4]{85}$ C. 16 D. $\sqrt[4]{75}$

9. $(4\sqrt{2})^2 =$

- A. 32 B. $16\sqrt{2}$ C. 64 D. 8

10. $(5 - \sqrt{7})(7 + \sqrt{7}) =$

- A. $42 - 2\sqrt{7}$ B. $28 - 2\sqrt{7}$ C. 28 D. $28 + 2\sqrt{7}$

11. A rationalising factor of $4\sqrt{10}$ is

- A. $\sqrt{10}$ B. 4 C. $4 + \sqrt{10}$ D. 10

12. $(\sqrt[3]{11} - \sqrt[3]{4})((\sqrt[3]{11})^2 + \sqrt[3]{11} \cdot \sqrt[3]{4} + (\sqrt[3]{4})^2) =$

A. 1267

B. 15

C. 7

D. $\sqrt[3]{7}$

13. $\sqrt{175} + \sqrt{252} =$

A. $\sqrt{427}$

B. $30\sqrt{7}$

C. $11\sqrt{14}$

D. $11\sqrt{7}$

14. $\sqrt{28}$ in simplest form is

A. $4\sqrt{7}$

B. $7\sqrt{2}$

C. $2\sqrt{7}$

D. $2\sqrt{14}$

15. $(\sqrt{2})^{13} =$

A. $64\sqrt{2}$

B. 64

C. $32\sqrt{2}$

D. $128\sqrt{2}$

16. $(\sqrt{10} - \sqrt{2})^2 =$

A. $12 - 4\sqrt{5}$

B. 8

C. $104 - 4\sqrt{5}$

D. $12 + 4\sqrt{5}$

17. $(\sqrt{6} - \sqrt{3})(\sqrt{6} + \sqrt{3}) =$

A. 9

B. 27

C. 3

D. $\sqrt{3}$

18. $\frac{1}{\sqrt{17}-\sqrt{10}} =$

A. $\frac{\sqrt{17}+\sqrt{10}}{27}$

B. $\sqrt{17} + \sqrt{10}$

C. $\frac{\sqrt{17}-\sqrt{10}}{7}$

D. $\frac{\sqrt{17}+\sqrt{10}}{7}$

19. $\frac{1}{7+\sqrt{13}} =$

A. $\frac{7-\sqrt{13}}{62}$

B. $\frac{\sqrt{13}-7}{6}$

C. $\frac{7+\sqrt{13}}{36}$

D. $\frac{7-\sqrt{13}}{36}$

20. If $\frac{3+5\sqrt{3}}{3-\sqrt{3}} = a + b\sqrt{3}$, then $b =$

A. 12

B. 4

C. 3

D. -3

21. $\frac{11+\sqrt{17}}{11-\sqrt{17}} + \frac{11-\sqrt{17}}{11+\sqrt{17}} =$

A. $\frac{69}{52}$

B. $-\frac{69}{26}$

C. $\frac{11\sqrt{17}}{26}$

D. $\frac{69}{26}$

22. $\frac{1}{8+\sqrt{6}} + \frac{1}{8-\sqrt{6}} =$

A. $\frac{8}{29}$

B. $\frac{\sqrt{6}}{29}$

C. 0

D. $\frac{8}{35}$

23. If $x = \sqrt[3]{13 + 2\sqrt{42}}$, then $x^3 + \frac{1}{x^3} =$

A. $4\sqrt{42}$

B. 13

C. 0

D. 26

24. $\sqrt{15 + 10\sqrt{2}} =$

A. $\sqrt{10} - \sqrt{5}$

B. $\sqrt{50} + 1$

C. $\sqrt{10} + \sqrt{5}$

D. $\sqrt{5} - \sqrt{10}$

25. $\sqrt{16 - \sqrt{87}} =$

A. $\frac{\sqrt{29}-\sqrt{3}}{\sqrt{2}}$

B. $\frac{\sqrt{29}-\sqrt{3}}{2}$

C. $\sqrt{29} - \sqrt{3}$

D. $\frac{\sqrt{29}+\sqrt{3}}{\sqrt{2}}$

Key sheet

1. A	2. B	3. D	4. B	5. B
6. C	7. B	8. A	9. A	10. B
11. A	12. C	13. D	14. C	15. A
16. A	17. C	18. D	19. D	20. C
21. D	22. A	23. D	24. C	25. A

Solutions

1. Which of the following is a rational number?

$\sqrt[3]{64} = \sqrt[3]{4 \times 4 \times 4} = 4$, a rational number, so it is not a surd.

The other three are roots of numbers that are not perfect powers, so they are surds.

Ans: (A) $\sqrt[3]{64}$

2. $\sqrt[6]{15}$ and e are both irrational. Which of them is a surd?

$1^6 = 1$ and $2^6 = 64$, so 15 is not a perfect sixth power. So $\sqrt[6]{15}$ is a surd.

e is irrational but it is not a root of a rational number, so it is not a surd.

Ans: (B) $\sqrt[6]{15}$ only

3. $9\sqrt[3]{n}$ is a mixed surd when n is

$2^3 = 8$ and $3^3 = 27$, so 19 is not a perfect cube. So $9\sqrt[3]{19}$ is a mixed surd.

For the others $\sqrt[3]{n}$ is a whole number, so the product is rational.

Ans: (D) 19

4. The conjugate surd of $12 - \sqrt{30}$ is

Only the sign of the surd part changes: the conjugate of $12 - \sqrt{30}$ is $12 + \sqrt{30}$.

Ans: (B) $12 + \sqrt{30}$

5. $\sqrt{10} \times \sqrt{10} =$

$$\sqrt{10} \times \sqrt{10} = \sqrt{10^2} = 10$$

Ans: (B) 10

6. $\frac{\sqrt{171}}{\sqrt{19}} =$

$$\begin{aligned}\frac{\sqrt{171}}{\sqrt{19}} &= \sqrt{\frac{171}{19}} \\ &= \sqrt{9} = \sqrt{3 \times 3} = 3\end{aligned}$$

Ans: (C) 3

7. $64^{\frac{1}{6}} =$

$$64^{\frac{1}{6}} = \sqrt[6]{64} = \sqrt[6]{2^6} = 2$$

Ans: (B) 2

8. $\frac{\sqrt[4]{80}}{\sqrt[4]{5}} =$

$$\begin{aligned}\frac{\sqrt[4]{80}}{\sqrt[4]{5}} &= \sqrt[4]{\frac{80}{5}} \\ &= \sqrt[4]{16} = \sqrt[4]{2^4} = 2\end{aligned}$$

Ans: (A) 2

9. $(4\sqrt{2})^2 =$

$$\begin{aligned}(4\sqrt{2})^2 &= 4^2 \times (\sqrt{2})^2 \\ &= 16 \times 2 = 32\end{aligned}$$

Ans: (A) 32

10. $(5 - \sqrt{7})(7 + \sqrt{7}) =$

$$(5 - \sqrt{7})(7 + \sqrt{7}) = 35 + 5\sqrt{7} - 7\sqrt{7} - 7$$

$$= 28 - 2\sqrt{7}$$

Ans: (B) $28 - 2\sqrt{7}$

11. A rationalising factor of $4\sqrt{10}$ is

$$4\sqrt{10} \times \sqrt{10} = 4 \times 10 = 40, \text{ a rational number.}$$

Ans: (A) $\sqrt{10}$

12. $(\sqrt[3]{11} - \sqrt[3]{4})((\sqrt[3]{11})^2 + \sqrt[3]{11} \cdot \sqrt[3]{4} + (\sqrt[3]{4})^2) =$

Use $(x - y)(x^2 + xy + y^2) = x^3 - y^3$ with $x = \sqrt[3]{11}$, $y = \sqrt[3]{4}$.

$$\text{Product} = (\sqrt[3]{11})^3 - (\sqrt[3]{4})^3 = 11 - 4 = 7$$

Ans: (C) 7

13. $\sqrt{175} + \sqrt{252} =$

$$\sqrt{175} = \sqrt{25 \times 7} = 5\sqrt{7}$$

$$\sqrt{252} = \sqrt{36 \times 7} = 6\sqrt{7}$$

$$5\sqrt{7} + 6\sqrt{7} = 11\sqrt{7} \text{ (add like surds)}$$

Ans: (D) $11\sqrt{7}$

14. $\sqrt{28}$ in simplest form is

$$\sqrt{28} = \sqrt{4 \times 7} = 2\sqrt{7}$$

Ans: (C) $2\sqrt{7}$

15. $(\sqrt{2})^{13} =$

$$(\sqrt{2})^{13} = [(\sqrt{2})^2]^6 \times \sqrt{2} = 2^6 \times \sqrt{2} = 64\sqrt{2}$$

Ans: (A) $64\sqrt{2}$

16. $(\sqrt{10} - \sqrt{2})^2 =$

$$(\sqrt{10} - \sqrt{2})^2 = 10 + 2 - 2\sqrt{20} = 12 - 4\sqrt{5}$$

Ans: (A) $12 - 4\sqrt{5}$

17. $(\sqrt{6} - \sqrt{3})(\sqrt{6} + \sqrt{3}) =$

$$(a + b)(a - b) = a^2 - b^2$$

$$= (\sqrt{6})^2 - (\sqrt{3})^2 = 6 - 3 = 3$$

Ans: (C) 3

18. $\frac{1}{\sqrt{17} - \sqrt{10}} =$

Multiply above and below by $\sqrt{17} + \sqrt{10}$: $(\sqrt{17} - \sqrt{10})(\sqrt{17} + \sqrt{10}) = 17 - 10 = 7$

$$= \frac{\sqrt{17} + \sqrt{10}}{7}$$

Ans: (D) $\frac{\sqrt{17} + \sqrt{10}}{7}$

19. $\frac{1}{7+\sqrt{13}} =$

Multiply above and below by $7 - \sqrt{13}$: $(7 + \sqrt{13})(7 - \sqrt{13}) = 49 - 13 = 36$

$$= \frac{7-\sqrt{13}}{36}$$

Ans: (D) $\frac{7-\sqrt{13}}{36}$

20. If $\frac{3+5\sqrt{3}}{3-\sqrt{3}} = a + b\sqrt{3}$, then $b =$

$$\frac{3+5\sqrt{3}}{3-\sqrt{3}} = \frac{(3+5\sqrt{3})(3+\sqrt{3})}{(3-\sqrt{3})(3+\sqrt{3})}$$

Numerator: $9 + 3\sqrt{3} + 15\sqrt{3} + 15 = 24 + 18\sqrt{3}$

Denominator: $3^2 - (\sqrt{3})^2 = 9 - 3 = 6$

$$\frac{24+18\sqrt{3}}{6} = 4 + 3\sqrt{3}$$

So $a = 4, b = 3$

Ans: (C) 3

21. $\frac{11+\sqrt{17}}{11-\sqrt{17}} + \frac{11-\sqrt{17}}{11+\sqrt{17}} =$

$$= \frac{(11+\sqrt{17})^2 + (11-\sqrt{17})^2}{(11+\sqrt{17})(11-\sqrt{17})} = \frac{276}{104} = \frac{69}{26}$$

The $\sqrt{17}$ terms cancel and $(11 + \sqrt{17})(11 - \sqrt{17}) = 104$.

Ans: (D) $\frac{69}{26}$

22. $\frac{1}{8+\sqrt{6}} + \frac{1}{8-\sqrt{6}} =$

$$= \frac{(8-\sqrt{6})+(8+\sqrt{6})}{(8+\sqrt{6})(8-\sqrt{6})} = \frac{16}{58} = \frac{8}{29}$$

Ans: (A) $\frac{8}{29}$

23. If $x = \sqrt[3]{13 + 2\sqrt{42}}$, then $x^3 + \frac{1}{x^3} =$

$$x^3 = 13 + 2\sqrt{42}, \text{ so } \frac{1}{x^3} = \frac{1}{13+2\sqrt{42}} = 13 - 2\sqrt{42}$$

$$x^3 + \frac{1}{x^3} = (13 + 2\sqrt{42}) + (13 - 2\sqrt{42}) = 26$$

Ans: (D) 26

24. $\sqrt{15 + 10\sqrt{2}} =$

$$10\sqrt{2} = 2 \times 5\sqrt{2} = 2\sqrt{50}$$

$$\sqrt{15 + 10\sqrt{2}} = \sqrt{10 + 5 + 2\sqrt{10 \times 5}}$$

$$= \sqrt{(\sqrt{10})^2 + (\sqrt{5})^2 + 2(\sqrt{10})(\sqrt{5})}$$

$$= \sqrt{(\sqrt{10} + \sqrt{5})^2} = \sqrt{10} + \sqrt{5}$$

Ans: (C) $\sqrt{10} + \sqrt{5}$

25. $\sqrt{16 - \sqrt{87}} =$

$$\sqrt{16 - \sqrt{87}} = \sqrt{\frac{2(16-\sqrt{87})}{2}} = \frac{\sqrt{32-2\sqrt{87}}}{\sqrt{2}}$$

$$= \frac{\sqrt{29+3-2\sqrt{29 \times 3}}}{\sqrt{2}}$$

$$= \frac{\sqrt{(\sqrt{29})^2 + (\sqrt{3})^2 - 2(\sqrt{29})(\sqrt{3})}}{\sqrt{2}}$$

$$= \frac{\sqrt{(\sqrt{29}-\sqrt{3})^2}}{\sqrt{2}} = \frac{\sqrt{29}-\sqrt{3}}{\sqrt{2}}$$

Ans: (A) $\frac{\sqrt{29}-\sqrt{3}}{\sqrt{2}}$