

Rationalization (Surds)

MCQ Test, Set 04

Time: 30 minutes

Maximum marks: 25

Name: _____ Class: _____ Date: _____

- There are 25 questions. Each has exactly one correct answer: choose A, B, C or D.
- Each correct answer carries 1 mark. There is no negative marking.
- The key sheet and worked solutions are at the end of this test.

1. For which value of n is $\sqrt[3]{n}$ a rational number?

- A. 37 B. 216 C. 33 D. 21

2. Which of the following statements is false?

- A. Every surd is an irrational number.
 B. Every irrational number is a surd.
 C. e is an irrational number.
 D. $\sqrt{48}$ is a surd.

3. Which one of these is a mixed surd?

- A. $9\sqrt{100}$ B. $\sqrt[4]{16}$ C. $\sqrt[3]{44}$ D. $5\sqrt[3]{29}$

4. The conjugate of $11 - \sqrt{21}$ is

- A. $-11 + \sqrt{21}$ B. $-11 - \sqrt{21}$ C. $11 + \sqrt{21}$ D. $\frac{1}{11 - \sqrt{21}}$

5. $\sqrt{6} \times \sqrt{24} =$

- A. 144 B. $6\sqrt{24}$ C. $\sqrt{30}$ D. 12

6. $\frac{\sqrt{192}}{\sqrt{3}} =$

- A. 64 B. $\sqrt{189}$ C. 8 D. $8\sqrt{3}$

7. $17^{\frac{1}{6}} =$

- A. $\frac{17}{6}$ B. $6\sqrt{17}$ C. $\sqrt[17]{6}$ D. $\sqrt[6]{17}$

8. $\frac{\sqrt[5]{729}}{\sqrt[5]{3}} =$

- A. 243 B. 3 C. $\sqrt[5]{726}$ D. $\sqrt[5]{732}$

9. $(2\sqrt{7})^2 =$

- A. 28 B. $4\sqrt{7}$ C. 14 D. 196

10. $(4 + \sqrt{5})(3 + \sqrt{5}) =$

- A. 17 B. $17 - 7\sqrt{5}$ C. $17 + 7\sqrt{5}$ D. $7 + 7\sqrt{5}$

11. A rationalising factor of $5\sqrt{14}$ is
A. $\sqrt{14}$ **B.** 14 **C.** 5 **D.** $5 + \sqrt{14}$
-
12. $(\sqrt[3]{13} - \sqrt[3]{11})((\sqrt[3]{13})^2 + \sqrt[3]{13} \cdot \sqrt[3]{11} + (\sqrt[3]{11})^2) =$
A. $\sqrt[3]{2}$ **B.** 24 **C.** 2 **D.** 866
-
13. $\sqrt{200}$ in simplest form is
A. $10\sqrt{2}$ **B.** $10\sqrt{20}$ **C.** $2\sqrt{10}$ **D.** $100\sqrt{2}$
-
14. $(\sqrt{15})^4 =$
A. $15\sqrt{15}$ **B.** 225 **C.** 60 **D.** 50625
-
15. $(\sqrt{11} + \sqrt{10})^2 =$
A. $21 + 2\sqrt{110}$ **B.** $21 - 2\sqrt{110}$ **C.** $221 + 2\sqrt{110}$ **D.** 21
-
16. $(\sqrt{19} - \sqrt{14})(\sqrt{19} + \sqrt{14}) =$
A. 165 **B.** $\sqrt{5}$ **C.** 5 **D.** 33
-
17. $(3 + 5\sqrt{7})(3 - 5\sqrt{7}) =$
A. 184 **B.** -26 **C.** -166 **D.** -1216
-
18. $\frac{1}{9+\sqrt{11}} =$
A. $\frac{\sqrt{11}-9}{2}$ **B.** $\frac{9-\sqrt{11}}{70}$ **C.** $\frac{9+\sqrt{11}}{70}$ **D.** $\frac{9-\sqrt{11}}{92}$
-
19. If $\frac{6+3\sqrt{7}}{4-\sqrt{7}} = a + b\sqrt{7}$, then $a =$
A. 45 **B.** 2 **C.** -5 **D.** 5
-
20. $\frac{\sqrt{19}+\sqrt{5}}{\sqrt{19}-\sqrt{5}} + \frac{\sqrt{19}-\sqrt{5}}{\sqrt{19}+\sqrt{5}} =$
A. $-\frac{24}{7}$ **B.** $\frac{2\sqrt{95}}{7}$ **C.** $\frac{12}{7}$ **D.** $\frac{24}{7}$
-
21. $\frac{1}{\sqrt{7}+\sqrt{5}} + \frac{1}{\sqrt{7}-\sqrt{5}} =$
A. $\sqrt{5}$ **B.** 0 **C.** $\sqrt{7}$ **D.** $\frac{\sqrt{7}}{6}$
-
22. If $x = \sqrt[3]{15 - 4\sqrt{14}}$, then $x^3 + \frac{1}{x^3} =$
A. 30 **B.** 0 **C.** $8\sqrt{14}$ **D.** 15
-
23. $\sqrt{9 + 2\sqrt{18}} =$
A. $\sqrt{3} - \sqrt{6}$ **B.** $\sqrt{18} + 1$ **C.** $\sqrt{6} + \sqrt{3}$ **D.** $\sqrt{6} - \sqrt{3}$
-
24. $\sqrt{11 - 2\sqrt{30}} =$
A. $\sqrt{5} - \sqrt{6}$ **B.** $\sqrt{6} - \sqrt{5}$ **C.** $\sqrt{6} + \sqrt{5}$ **D.** $\sqrt{30} - 1$
-
25. $\sqrt{16 + \sqrt{207}} =$
A. $\sqrt{23} + 3$ **B.** $\frac{\sqrt{23}-3}{\sqrt{2}}$
C. $\frac{\sqrt{23}+3}{2}$ **D.** $\frac{\sqrt{23}+3}{\sqrt{2}}$
-

Key sheet

1. B	2. B	3. D	4. C	5. D
6. C	7. D	8. B	9. A	10. C
11. A	12. C	13. A	14. B	15. A
16. C	17. C	18. B	19. D	20. D
21. C	22. A	23. C	24. B	25. D

Solutions

1. For which value of n is $\sqrt[3]{n}$ a rational number?

$$\sqrt[3]{216} = \sqrt[3]{6 \times 6 \times 6} = 6$$

The other three are not perfect cubes, so their roots are surds.

Ans: (B) 216

2. Which of the following statements is false?

π and e are irrational numbers but not surds.

So 'every irrational number is a surd' is false.

Ans: (B) Every irrational number is a surd.

3. Which one of these is a mixed surd?

$5\sqrt[3]{29} = 5 \cdot \sqrt[3]{29}$: a rational number times a surd.

$\sqrt[3]{44}$ is a pure surd; $9\sqrt{100} = 90$ and $\sqrt[4]{16} = 2$ are rational.

Ans: (D) $5\sqrt[3]{29}$

4. The conjugate of $11 - \sqrt{21}$ is

Only the sign of the surd part changes: the conjugate of $11 - \sqrt{21}$ is $11 + \sqrt{21}$.

Ans: (C) $11 + \sqrt{21}$

5. $\sqrt{6} \times \sqrt{24} =$

$$\sqrt{6} \times \sqrt{24} = \sqrt{144} = \sqrt{12 \times 12} = 12$$

Ans: (D) 12

6. $\frac{\sqrt{192}}{\sqrt{3}} =$

$$\frac{\sqrt{192}}{\sqrt{3}} = \sqrt{\frac{192}{3}}$$

$$= \sqrt{64} = \sqrt{8 \times 8} = 8$$

Ans: (C) 8

7. $17^{\frac{1}{6}} =$

$$a^{\frac{1}{n}} = \sqrt[n]{a}, \text{ so } 17^{\frac{1}{6}} = \sqrt[6]{17}$$

Ans: (D) $\sqrt[6]{17}$

8. $\frac{\sqrt[5]{729}}{\sqrt[5]{3}} =$

$$\frac{\sqrt[5]{729}}{\sqrt[5]{3}} = \sqrt[5]{\frac{729}{3}}$$

$$= \sqrt[5]{243} = \sqrt[5]{3^5} = 3$$

Ans: (B) 3

9. $(2\sqrt{7})^2 =$

$$(2\sqrt{7})^2 = 2^2 \times (\sqrt{7})^2$$

$$= 4 \times 7 = 28$$

Ans: (A) 28

10. $(4 + \sqrt{5})(3 + \sqrt{5}) =$

$$(4 + \sqrt{5})(3 + \sqrt{5}) = 12 + 4\sqrt{5} + 3\sqrt{5} + 5 \\ = 17 + 7\sqrt{5}$$

Ans: (C) $17 + 7\sqrt{5}$

11. A rationalising factor of $5\sqrt{14}$ is

$$5\sqrt{14} \times \sqrt{14} = 5 \times 14 = 70, \text{ a rational number.}$$

Ans: (A) $\sqrt{14}$

12. $(\sqrt[3]{13} - \sqrt[3]{11})((\sqrt[3]{13})^2 + \sqrt[3]{13} \cdot \sqrt[3]{11} + (\sqrt[3]{11})^2) =$

Use $(x - y)(x^2 + xy + y^2) = x^3 - y^3$ with $x = \sqrt[3]{13}$, $y = \sqrt[3]{11}$.

$$\text{Product} = (\sqrt[3]{13})^3 - (\sqrt[3]{11})^3 = 13 - 11 = 2$$

Ans: (C) 2

13. $\sqrt{200}$ in simplest form is

$$\sqrt{200} = \sqrt{100 \times 2} = 10\sqrt{2}$$

Ans: (A) $10\sqrt{2}$

14. $(\sqrt{15})^4 =$

$$(\sqrt{15})^4 = [(\sqrt{15})^2]^2 = 15^2 = 225$$

Ans: (B) 225

15. $(\sqrt{11} + \sqrt{10})^2 =$

$$(\sqrt{11} + \sqrt{10})^2 = 11 + 10 + 2\sqrt{110} = 21 + 2\sqrt{110}$$

Ans: (A) $21 + 2\sqrt{110}$

16. $(\sqrt{19} - \sqrt{14})(\sqrt{19} + \sqrt{14}) =$

$$(a + b)(a - b) = a^2 - b^2$$

$$= (\sqrt{19})^2 - (\sqrt{14})^2 = 19 - 14 = 5$$

Ans: (C) 5

17. $(3 + 5\sqrt{7})(3 - 5\sqrt{7}) =$

$$(a + b)(a - b) = a^2 - b^2$$

$$= 3^2 - (5\sqrt{7})^2 = 9 - 175 = -166$$

Ans: (C) -166

18. $\frac{1}{9 + \sqrt{11}} =$

Multiply above and below by $9 - \sqrt{11}$: $(9 + \sqrt{11})(9 - \sqrt{11}) = 81 - 11 = 70$

$$= \frac{9 - \sqrt{11}}{70}$$

Ans: (B) $\frac{9 - \sqrt{11}}{70}$

19. If $\frac{6+3\sqrt{7}}{4-\sqrt{7}} = a + b\sqrt{7}$, then $a =$

$$\frac{6+3\sqrt{7}}{4-\sqrt{7}} = \frac{(6+3\sqrt{7})(4+\sqrt{7})}{(4-\sqrt{7})(4+\sqrt{7})}$$

$$\text{Numerator: } 24 + 6\sqrt{7} + 12\sqrt{7} + 21 = 45 + 18\sqrt{7}$$

$$\text{Denominator: } 4^2 - (\sqrt{7})^2 = 16 - 7 = 9$$

$$\frac{45+18\sqrt{7}}{9} = 5 + 2\sqrt{7}$$

So $a = 5, b = 2$

Ans: (D) 5

20. $\frac{\sqrt{19}+\sqrt{5}}{\sqrt{19}-\sqrt{5}} + \frac{\sqrt{19}-\sqrt{5}}{\sqrt{19}+\sqrt{5}} =$

$$= \frac{(\sqrt{19}+\sqrt{5})^2 + (\sqrt{19}-\sqrt{5})^2}{(\sqrt{19}+\sqrt{5})(\sqrt{19}-\sqrt{5})} = \frac{48}{14} = \frac{24}{7}$$

The $\sqrt{95}$ terms cancel and $(\sqrt{19} + \sqrt{5})(\sqrt{19} - \sqrt{5}) = 19 - 5 = 14$.

Ans: (D) $\frac{24}{7}$

21. $\frac{1}{\sqrt{7}+\sqrt{5}} + \frac{1}{\sqrt{7}-\sqrt{5}} =$

$$= \frac{(\sqrt{7}-\sqrt{5}) + (\sqrt{7}+\sqrt{5})}{(\sqrt{7}+\sqrt{5})(\sqrt{7}-\sqrt{5})} = \frac{2\sqrt{7}}{2} = \sqrt{7}$$

Ans: (C) $\sqrt{7}$

22. If $x = \sqrt[3]{15 - 4\sqrt{14}}$, then $x^3 + \frac{1}{x^3} =$

$$x^3 = 15 - 4\sqrt{14}, \text{ so } \frac{1}{x^3} = \frac{1}{15-4\sqrt{14}} = 15 + 4\sqrt{14}$$

$$x^3 + \frac{1}{x^3} = (15 - 4\sqrt{14}) + (15 + 4\sqrt{14}) = 30$$

Ans: (A) 30

23. $\sqrt{9 + 2\sqrt{18}} =$

$$\sqrt{9 + 2\sqrt{18}} = \sqrt{6 + 3 + 2\sqrt{6} \times 3}$$

$$= \sqrt{(\sqrt{6})^2 + (\sqrt{3})^2 + 2(\sqrt{6})(\sqrt{3})}$$

$$= \sqrt{(\sqrt{6} + \sqrt{3})^2} = \sqrt{6} + \sqrt{3}$$

Ans: (C) $\sqrt{6} + \sqrt{3}$

24. $\sqrt{11 - 2\sqrt{30}} =$

$$\sqrt{11 - 2\sqrt{30}} = \sqrt{6 + 5 - 2\sqrt{6} \times 5}$$

$$= \sqrt{(\sqrt{6})^2 + (\sqrt{5})^2 - 2(\sqrt{6})(\sqrt{5})}$$

$$= \sqrt{(\sqrt{6} - \sqrt{5})^2} = \sqrt{6} - \sqrt{5}$$

Ans: (B) $\sqrt{6} - \sqrt{5}$

25. $\sqrt{16 + \sqrt{207}} =$

$$\begin{aligned}\sqrt{16 + \sqrt{207}} &= \sqrt{\frac{2(16 + \sqrt{207})}{2}} = \frac{\sqrt{32 + 2\sqrt{207}}}{\sqrt{2}} \\&= \frac{\sqrt{23 + 9 + 2\sqrt{23 \times 9}}}{\sqrt{2}} \\&= \frac{\sqrt{(\sqrt{23})^2 + (3)^2 + 2(\sqrt{23})(3)}}{\sqrt{2}} \\&= \frac{\sqrt{(\sqrt{23} + 3)^2}}{\sqrt{2}} = \frac{\sqrt{23} + 3}{\sqrt{2}}\end{aligned}$$

Ans: (D) $\frac{\sqrt{23} + 3}{\sqrt{2}}$
