

# Rationalization (Surds)

## Question Paper, Set 01

Time: 1 hour 30 minutes

Maximum marks: 50

Name: \_\_\_\_\_ Class: \_\_\_\_\_ Date: \_\_\_\_\_

1. Answer all 20 questions. The marks for each question are shown in brackets.
2. Show your working for the 2, 3 and 5 mark questions.
3. The key sheet and worked solutions are at the end of this paper.

1. Is  $\sqrt[6]{10}$  a surd? Give a reason. [1 mark]
2. Is  $2\sqrt{49}$  a mixed surd? Give a reason. [1 mark]
3. Write the conjugates of (i)  $9 - \sqrt{5}$  (ii)  $\sqrt{15} + 5$  [2 marks]
4. Find  $\sqrt{38} \times \sqrt{38}$ . [1 mark]
5. Find the value of  $625^{\frac{1}{2}}$ . [2 marks]
6. Simplify (i)  $\sqrt[6]{16} \times \sqrt[6]{4}$  (ii)  $\frac{\sqrt[3]{243}}{\sqrt[3]{9}}$  [3 marks]
7. Simplify  $(12\sqrt{3})^2$ . [1 mark]
8. Simplify  $(1 + \sqrt{11})(3 + \sqrt{11})$ . [2 marks]
9. Write the rationalising factor of  $11 - \sqrt{26}$ . [1 mark]
10. Simplify  $\sqrt{216} - \sqrt{54}$ . [2 marks]
11. Find (i)  $(\sqrt{6})^5$  (ii)  $(4\sqrt{10})^3$  [3 marks]
12. Simplify  $(10 + \sqrt{11})^2$ . [2 marks]
13. Find  $(19 + \sqrt{19})(19 - \sqrt{19})$ . [1 mark]
14. Rationalise the denominator of  $\frac{1}{4 + \sqrt{5}}$ . [2 marks]
15. If  $\frac{\sqrt{6} + \sqrt{5}}{\sqrt{6} - \sqrt{5}} = a + b\sqrt{30}$ , find  $a$  and  $b$ . [5 marks]
16. Simplify  $\frac{3 + \sqrt{3}}{3 - \sqrt{3}} + \frac{3 - \sqrt{3}}{3 + \sqrt{3}}$ . [3 marks]
17. Simplify  $\frac{4}{\sqrt{10} - \sqrt{6}} + \frac{6}{2 - \sqrt{10}} - \frac{2}{\sqrt{6} + 2}$ . [5 marks]
18. If  $x = \frac{1}{3 + 2\sqrt{2}}$ , find the value of  $x^3 - 7x^2 + 7x + 4$ . [5 marks]
19. Find  $\sqrt{7 + \sqrt{24}}$ . [3 marks]

**20.** Find  $\sqrt{13 - \sqrt{69}}$  and write the answer with a rational denominator.

---

[5 marks]

## Key sheet

| Q  | Answer                                                                   |
|----|--------------------------------------------------------------------------|
| 1  | Yes. 10 is not a perfect sixth power, so $\sqrt[6]{10}$ is not rational. |
| 2  | No. $2\sqrt{49} = 14$ , a rational number.                               |
| 3  | (i) $9 + \sqrt{5}$ (ii) $-\sqrt{15} + 5$                                 |
| 4  | 38                                                                       |
| 5  | 25                                                                       |
| 6  | (i) 2 (ii) 3                                                             |
| 7  | 432                                                                      |
| 8  | $14 + 4\sqrt{11}$                                                        |
| 9  | $11 + \sqrt{26}$                                                         |
| 10 | $3\sqrt{6}$                                                              |
| 11 | (i) $36\sqrt{6}$ (ii) $640\sqrt{10}$                                     |
| 12 | $111 + 20\sqrt{11}$                                                      |
| 13 | 342                                                                      |
| 14 | $\frac{4-\sqrt{5}}{11}$                                                  |
| 15 | $a = 11, b = 2$                                                          |
| 16 | 4                                                                        |
| 17 | 0                                                                        |
| 18 | 5                                                                        |
| 19 | $\sqrt{6} + 1$                                                           |
| 20 | $\frac{\sqrt{46}-\sqrt{6}}{2}$                                           |

## Solutions

---

1. Is  $\sqrt[6]{10}$  a surd? Give a reason. [1]

$1^6 = 1$  and  $2^6 = 64$ , so 10 is not a perfect sixth power.

So  $\sqrt[6]{10}$  is not a rational number; it is a surd.

**Ans: Yes. 10 is not a perfect sixth power, so  $\sqrt[6]{10}$  is not rational.**

---

2. Is  $2\sqrt{49}$  a mixed surd? Give a reason. [1]

$$\sqrt{49} = \sqrt{7 \times 7} = 7$$

$2\sqrt{49} = 2 \times 7 = 14$ , which is rational, so it is not a surd.

**Ans: No.  $2\sqrt{49} = 14$ , a rational number.**

---

3. Write the conjugates of (i)  $9 - \sqrt{5}$  (ii)  $\sqrt{15} + 5$  [2]

Change the sign of the surd part only.

(i) The conjugate of  $9 - \sqrt{5}$  is  $9 + \sqrt{5}$

(ii) The conjugate of  $\sqrt{15} + 5$  is  $-\sqrt{15} + 5$

**Ans: (i)  $9 + \sqrt{5}$  (ii)  $-\sqrt{15} + 5$**

---

4. Find  $\sqrt{38} \times \sqrt{38}$ . [1]

$$\sqrt{38} \times \sqrt{38} = (\sqrt{38})^2$$

$$= \sqrt{38^2} = 38$$

**Ans: 38**

---

5. Find the value of  $625^{\frac{1}{2}}$ . [2]

$$625^{\frac{1}{2}} = \sqrt{625}$$

$$= \sqrt{25^2} = \sqrt{25 \times 25} = 25$$

**Ans: 25**

---

6. Simplify (i)  $\sqrt[6]{16} \times \sqrt[6]{4}$  (ii)  $\frac{\sqrt[3]{243}}{\sqrt[3]{9}}$  [3]

$$(i) \sqrt[6]{16} \times \sqrt[6]{4} = \sqrt[6]{16 \times 4}$$

$$= \sqrt[6]{64} = \sqrt[6]{2^6} = 2$$

$$(ii) \frac{\sqrt[3]{243}}{\sqrt[3]{9}} = \sqrt[3]{\frac{243}{9}}$$

$$= \sqrt[3]{27} = \sqrt[3]{3^3} = 3$$

**Ans: (i) 2 (ii) 3**

---

7. Simplify  $(12\sqrt{3})^2$ . [1]

$$(12\sqrt{3})^2 = 12^2 \times (\sqrt{3})^2$$

$$= 144 \times 3 = 432$$

**Ans: 432**

---

8. Simplify  $(1 + \sqrt{11})(3 + \sqrt{11})$ . [2]

$$(1 + \sqrt{11})(3 + \sqrt{11}) = 3 + \sqrt{11} + 3\sqrt{11} + 11 \\ = 14 + 4\sqrt{11}$$

**Ans:**  $14 + 4\sqrt{11}$

---

9. Write the rationalising factor of  $11 - \sqrt{26}$ . [1]

$$(11 - \sqrt{26})(11 + \sqrt{26}) = 11^2 - (\sqrt{26})^2 = 121 - 26 = 95, \text{ a rational number.}$$

So the rationalising factor is  $11 + \sqrt{26}$ .

**Ans:**  $11 + \sqrt{26}$

---

10. Simplify  $\sqrt{216} - \sqrt{54}$ . [2]

$$\sqrt{216} = \sqrt{36 \times 6} = 6\sqrt{6}$$

$$\sqrt{54} = \sqrt{9 \times 6} = 3\sqrt{6}$$

$$\sqrt{216} - \sqrt{54} = 6\sqrt{6} - 3\sqrt{6} = 3\sqrt{6}$$

**Ans:**  $3\sqrt{6}$

---

11. Find (i)  $(\sqrt{6})^5$  (ii)  $(4\sqrt{10})^3$  [3]

$$(i) (\sqrt{6})^5 = [(\sqrt{6})^2]^2 \times \sqrt{6} = 6^2 \times \sqrt{6} = 36\sqrt{6}$$

$$(ii) (4\sqrt{10})^3 = 4^3 \times (\sqrt{10})^3 = 64 \times 10\sqrt{10} = 640\sqrt{10}$$

**Ans:** (i)  $36\sqrt{6}$  (ii)  $640\sqrt{10}$

---

12. Simplify  $(10 + \sqrt{11})^2$ . [2]

$$(a + b)^2 = a^2 + b^2 + 2ab$$

$$(10 + \sqrt{11})^2 = 10^2 + (\sqrt{11})^2 + 2(10)(\sqrt{11})$$

$$= 100 + 11 + 20\sqrt{11} = 111 + 20\sqrt{11}$$

**Ans:**  $111 + 20\sqrt{11}$

---

13. Find  $(19 + \sqrt{19})(19 - \sqrt{19})$ . [1]

$$(a + b)(a - b) = a^2 - b^2$$

$$= 19^2 - (\sqrt{19})^2 = 361 - 19 = 342$$

**Ans:** 342

---

14. Rationalise the denominator of  $\frac{1}{4+\sqrt{5}}$ . [2]

$$\frac{1}{4+\sqrt{5}} = \frac{4-\sqrt{5}}{(4+\sqrt{5})(4-\sqrt{5})}$$

$$(4 + \sqrt{5})(4 - \sqrt{5}) = 4^2 - (\sqrt{5})^2 = 16 - 5 = 11$$

$$= \frac{4-\sqrt{5}}{11}$$

**Ans:**  $\frac{4-\sqrt{5}}{11}$

---

15. If  $\frac{\sqrt{6}+\sqrt{5}}{\sqrt{6}-\sqrt{5}} = a + b\sqrt{30}$ , find  $a$  and  $b$ . [5]

$$\begin{aligned}\frac{\sqrt{6}+\sqrt{5}}{\sqrt{6}-\sqrt{5}} &= \frac{(\sqrt{6}+\sqrt{5})^2}{(\sqrt{6}-\sqrt{5})(\sqrt{6}+\sqrt{5})} \\ &= \frac{6+5+2\sqrt{30}}{6-5} \\ &= 11 + 2\sqrt{30}\end{aligned}$$

Comparing with  $a + b\sqrt{30}$ :  $a = 11, b = 2$

Ans:  $a = 11, b = 2$

16. Simplify  $\frac{3+\sqrt{3}}{3-\sqrt{3}} + \frac{3-\sqrt{3}}{3+\sqrt{3}}$ . [3]

$$\begin{aligned}&= \frac{(3+\sqrt{3})^2}{(3-\sqrt{3})(3+\sqrt{3})} + \frac{(3-\sqrt{3})^2}{(3+\sqrt{3})(3-\sqrt{3})} \\ (3 + \sqrt{3})^2 &= 9 + 3 + 6\sqrt{3} = 12 + 6\sqrt{3}, (3 - \sqrt{3})^2 = 12 - 6\sqrt{3}, (3 + \sqrt{3})(3 - \sqrt{3}) = 6 \\ &= \frac{12+6\sqrt{3}}{6} + \frac{12-6\sqrt{3}}{6} \\ &= \frac{12+6\sqrt{3}+12-6\sqrt{3}}{6} = 4\end{aligned}$$

Ans: 4

17. Simplify  $\frac{4}{\sqrt{10}-\sqrt{6}} + \frac{6}{2-\sqrt{10}} - \frac{2}{\sqrt{6}+2}$ . [5]

$$\begin{aligned}\frac{4}{\sqrt{10}-\sqrt{6}} &= \frac{4(\sqrt{10}+\sqrt{6})}{(\sqrt{10}-\sqrt{6})(\sqrt{10}+\sqrt{6})} = \frac{4(\sqrt{10}+\sqrt{6})}{4} = \sqrt{10} + \sqrt{6} \\ \frac{6}{2-\sqrt{10}} &= \frac{6(2+\sqrt{10})}{(2-\sqrt{10})(2+\sqrt{10})} = \frac{6(2+\sqrt{10})}{-6} = -2 - \sqrt{10} \\ \frac{2}{\sqrt{6}+2} &= \frac{2(\sqrt{6}-2)}{(\sqrt{6}+2)(\sqrt{6}-2)} = \frac{2(\sqrt{6}-2)}{2} = \sqrt{6} - 2\end{aligned}$$

$$\text{Sum} = (\sqrt{10} + \sqrt{6}) + (-2 - \sqrt{10}) - (\sqrt{6} - 2)$$

= 0 (all the terms cancel)

Ans: 0

18. If  $x = \frac{1}{3+2\sqrt{2}}$ , find the value of  $x^3 - 7x^2 + 7x + 4$ . [5]

$$\begin{aligned}x &= \frac{1}{3+2\sqrt{2}} = \frac{3-2\sqrt{2}}{(3+2\sqrt{2})(3-2\sqrt{2})} \\ &= \frac{3-2\sqrt{2}}{9-8} = 3 - 2\sqrt{2}\end{aligned}$$

$$x - 3 = -2\sqrt{2}$$

$$(x - 3)^2 = (-2\sqrt{2})^2 \Rightarrow x^2 - 6x + 9 = 8$$

$$\Rightarrow x^2 - 6x + 1 = 0$$

Dividing  $x^3 - 7x^2 + 7x + 4$  by  $x^2 - 6x + 1$ : quotient  $x - 1$ , remainder 5

$$x^3 - 7x^2 + 7x + 4 = (x^2 - 6x + 1)(x - 1) + 5$$

$$= 0 \times (x - 1) + 5 = 5$$

Ans: 5

19. Find  $\sqrt{7 + \sqrt{24}}$ . [3]

$$\sqrt{24} = \sqrt{4 \times 6} = 2\sqrt{6}$$

$$\sqrt{7 + \sqrt{24}} = \sqrt{6 + 1 + 2\sqrt{6} \times 1}$$

$$= \sqrt{(\sqrt{6})^2 + (1)^2 + 2(\sqrt{6})(1)}$$

$$= \sqrt{(\sqrt{6} + 1)^2} = \sqrt{6} + 1$$

Ans:  $\sqrt{6} + 1$

20. Find  $\sqrt{13 - \sqrt{69}}$  and write the answer with a rational denominator. [5]

$$\begin{aligned}\sqrt{13 - \sqrt{69}} &= \sqrt{\frac{2(13 - \sqrt{69})}{2}} = \frac{\sqrt{26 - 2\sqrt{69}}}{\sqrt{2}} \\&= \frac{\sqrt{23+3-2\sqrt{23 \times 3}}}{\sqrt{2}} \\&= \frac{\sqrt{(\sqrt{23})^2 + (\sqrt{3})^2 - 2(\sqrt{23})(\sqrt{3})}}{\sqrt{2}} \\&= \frac{\sqrt{(\sqrt{23} - \sqrt{3})^2}}{\sqrt{2}} = \frac{\sqrt{23} - \sqrt{3}}{\sqrt{2}} \\&= \frac{\sqrt{23} - \sqrt{3}}{\sqrt{2}} = \frac{(\sqrt{23} - \sqrt{3}) \times \sqrt{2}}{\sqrt{2} \times \sqrt{2}} = \frac{\sqrt{46} - \sqrt{6}}{2}\end{aligned}$$

Ans:  $\frac{\sqrt{46} - \sqrt{6}}{2}$

---